

CHAPTER 32

Inflammation

Helene F. Rosenberg and John I. Gallin

Historical Perspective and Overview

Initiation of the Acute Inflammatory Response

Physical Responses to Acute Injury

Vasodilation • Increased Vascular Permeability • Neutrophil Recruitment and Activation • Fever

Molecular Mediators of the Acute Inflammatory Response

The Plasma Proteases • Lipid Mediators • Peptides and Amines • Nitric Oxide • Proinflammatory Cytokines

Model Systems of Inflammation

Response to Intravenous Endotoxin • Temporal Analysis of Soluble Mediators in Blister Fluid

Cellular Mediators of the Acute Inflammatory Response

Neutrophils • Monocytes and Macrophages • Eosinophils • Platelets and Lymphocytes • Endothelial Cells

Allergy and Inflammation

Resolution of the Acute Inflammatory Response

Cell Senescence or Apoptosis • Antiinflammatory Mediators • Hypothalamo-Pituitary-Adrenocortical Axis • Wound Repair and Angiogenesis

Chronic Inflammation

Future Directions: Novel Antiinflammatory Therapies

Immunophilin-binding Agents • Experimental Antiinflammatory Agents

References

HISTORICAL PERSPECTIVE AND OVERVIEW

Inflammation is the physiologic process by which vascularized tissues respond to injury. During the inflammatory process, soluble mediators and cellular components work together in a systematic fashion in the attempt to contain and eliminate the agents causing physical distress. While inflammation is clearly crucial to maintaining the health and integrity of an individual organism, when poorly controlled, the inflammatory process can result in massive tissue destruction. For this reason, the concept of inflammation as a double-edged sword has taken hold.

The first observations on the inflammatory process are credited to Cornelius Celsus, who described the cardinal clinical signs of inflammation during the first century of the Common Era. His signs—*rubor* (redness), *dolor* (pain), *calor* (heat), and *tumor* (swelling)—remain as focal points for study. Another early contributor to this field was John Hunter (1793), who was the first to appreciate inflammation as host defense, as opposed to a disease process (1). In the 1800s, Julius Cohnheim provided the first microscopic descriptions of the inflammatory process (2). Paul Ehrlich contributed to the overall understanding of the inflamma-

tory process with his observations on the role of antibodies, and Elie Metchnikoff, with his observations on phagocytosis; both were awarded the Nobel prize for their work in 1908. Other crucial discoveries included those of Wright, who described the plasma proteins (opsonins) that coat and tag foreign substances for phagocytic destruction, and Dale and Laidlaw, who demonstrated the vasoactive role of histamine (3). In recent history, many investigators have contributed observations on the soluble mediators known as cytokines (which include chemokines, interleukins, interferons, and colony-stimulating factors) and the role played by cytokines and their specific receptors in modulating nearly every event characteristic of the inflammatory response.

Inflammation has been traditionally divided into acute and chronic responses. Acute inflammation is the rapid, short-lived (minutes to days), relatively uniform response to acute injury, characterized by accumulations of fluid, plasma proteins, and neutrophilic leukocytes. In contrast, chronic inflammation is of longer duration and includes influx of lymphocytes and macrophages and fibroblast growth.

The highlights of the events characteristic of acute and chronic inflammation are as depicted in Fig. 1. For more detail on all topics covered in this chapter, the reader is referred to the textbook entitled, *Inflammation* (4).

An injuring agent evades or destroys primary barriers (epithelial or endothelial cells and their specialized structures), initiating

H. F. Rosenberg and J. I. Gallin: Laboratory of Host Defenses, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland 20892-1504.




FIG. 1. Molecular and cellular events of the inflammatory response.

acute inflammation. Examples of injurious agents include pathogens (e.g., bacteria, viruses, parasites), foreign bodies from exogenous (e.g., asbestos) or endogenous (e.g., urate crystals, immune complexes) sources, as well as physical (e.g., burns) or chemical (e.g., caustics) agents.

Tissue damage initiates a series of molecular events, resulting in the production of soluble proinflammatory mediators that promote the hallmark physical signs of inflammation, including increased blood flow and vascular permeability, migration of leukocytes from the peripheral blood into the tissues, accumulation of these leukocytes at the inflammatory focus, and activation of the leukocytes to destroy and (if possible) eliminate the foreign substance. These soluble mediators include the plasma protease systems, lipid mediators, and proinflammatory peptides and cytokines. Addi-

tional mediators secreted by activated leukocytes can serve to prolong the inflammatory response by both direct and indirect means.

As the foreign threat is eliminated, antiinflammatory mediators permit the process to wind down, so as to avoid unnecessary and excessive damage to the tissues surrounding the inflammatory focus. If this acute process results in only incomplete destruction and/or elimination of the foreign substance, the inflammatory process persists and expands its repertoire of soluble mediators and cellular components, a process referred to as chronic inflammation.

This chapter describes in some detail the physical, cellular, and molecular events underlying acute and chronic inflammation. We describe several clinical disorders in which deficient or deranged inflammatory responses play a crucial role, and conclude with a discussion of novel therapeutic agents designed to combat the rav-

ages of an excessive inflammatory response. References have been selected to include recent reviews and textbook chapters that cover the individual topics in greater detail.

INITIATION OF THE ACUTE INFLAMMATORY RESPONSE

The way in which the inflammatory process is initiated depends in part on the nature and portal of entry of the foreign substance and, to some degree, the nature and circumstances of a particular individual. Pathogens can initiate inflammation by a number of distinct and idiosyncratic mechanisms, including activation of the plasma protease systems by interaction with degradation products of the bacterial cell walls and by secretion of toxins that can activate the inflammatory response directly (5). Injured cells can release degradation products that initiate one or more of the plasma protease cascades, and can upregulate expression of proinflammatory cytokines that augment the inflammatory process.

PHYSICAL RESPONSES TO ACUTE INJURY

Regardless of the initiating agent, the physiologic changes accompanying acute inflammation encompass four main features.

Vasodilation

Vasodilation (often preceded by a brief period of vasoconstriction) is one of the earliest physical responses to acute tissue injury. The arterioles are the first to be involved, followed by the capillary beds, resulting in a net increase in blood flow. The increased blood flow results in the characteristic heat and redness (calor and rubor) associated with foci of acute inflammation.

Increased Vascular Permeability

Under normal conditions, the vascular endothelial cells function as a semipermeable membrane, restricting the plasma proteins to the intravascular space. In response to inflammatory stimuli, endothelial cells lining the venules contract, widening the intercellular junctions to produce gaps, permitting passage of plasma proteins (tumor) (see ref. 6 for a more complete discussion). More severe injury is associated with endothelial cell necrosis and increased leakage of plasma proteins and blood cells.

Neutrophil Recruitment and Activation

One of the initial and most crucial responses to acute inflammation is the recruitment of leukocytes (primarily neutrophils) from the bloodstream (and ultimately from the bone marrow) to the focus of inflammatory activity. The first step observed in this process is margination, as the neutrophils appear to roll slowly along the periphery of the blood vessel. This is followed by a more definitive sticking, or adherence stage. Neutrophils then migrate into the tissue, traveling through the enlarged endothelial cell junctions and the basement membrane. Under the influence of soluble chemotactic agents (see below), neutrophils are targeted to the site of inflammation, where they collect to form an inflammatory exudate known as pus. At this site, the neutrophils ingest pathogenic material by a process known as phagocytosis, and detoxify and

digest this material by the actions of endogenous oxidants and proteolytic enzymes.

Fever

Fever remains the most poorly understood of the acute inflammatory responses. Agents producing fever, known as pyrogens, are released from leukocytes in response to specific stimuli, such as bacterial endotoxin. Pyrogens exert their actions via the temperature-regulating mechanism of the hypothalamus. A number of soluble proinflammatory mediators (discussed below and in Chapter 22) have been implicated in this process, including interleukin-1 (IL-1), tumor necrosis factor- α (TNF- α) and prostaglandins. The beneficial role of fever with respect to the acute inflammatory response remains a mystery.

MOLECULAR MEDIATORS OF THE ACUTE INFLAMMATORY RESPONSE

The physiologic features of the inflammatory process are initiated, regulated, and ultimately eliminated by the actions of numerous entities collectively termed *soluble inflammatory mediators*. Some of these mediators exist in inactive form and are activated by products of acute inflammation; others are synthesized and/or released from cellular sources, also in response to the products of acute inflammation, or by other soluble inflammatory mediators. Although presented as separate components, it is important to appreciate that these mediators perform their functions via intersecting, coordinately regulated, and mutually augmenting pathways.

The Plasma Proteases

Among the central components of the inflammatory response are the three interacting groups of plasma proteases. It is through the actions of plasma proteins as they convert one another from inactive to active forms that many of the major soluble mediators of inflammation are generated.

Complement

This group of plasma proteins was initially identified on the basis of their ability to complement the bactericidal activities of antibodies. At present, there are nine proteins known as participants in the complement cascade, described in detail in Chapter 29. By serial and sequential proteolytic cleavage, the complement proteins become activated and promote the inflammatory response by binding to foreign organisms and enhancing phagocytosis (components C3b and C4b), by functioning as agents increasing vascular permeability and as chemoattractants for inflammatory cells (C3a and C5a), and by creating lytic multiprotein complexes (C5b-9) (7-11). The classical pathway of complement is activated by antigen-bound antibodies of the IgM or IgG class. The first complement component, C1, undergoes autocatalytic cleavage to produce C1s, which in turn catalyzes a specific cleavage of C4 to C4b and C4a; C4b then binds to the target antigen, and C4a combines with C2a (product of the proteolytic cleavage of C2 by C1s) to create a protease specific for the component C3, and so on. The alternative pathway is utilized by other initiating agents (e.g., bacterial endo-

toxin) and converges with the classical pathway at the cleavage of C3; in addition, proteases from bacteria and damaged tissue, as well as plasmin generated by the fibrinolytic system (see below), can catalyze the cleavage of C3. From this point, the cleavage product C3b goes on to cleave C5, and serial proteolysis leads to activation of proteins C6 through C9.

Inherited deficiencies in individual components of the complement system can result in increased susceptibility to infection, rheumatic disorders, or angioedema. Individual deficiencies of various components have been identified (12,13); among the more common of these conditions is C2 deficiency. Individuals with C2 deficiency cannot utilize the classical pathway of complement activation; this autosomal recessive trait predisposes affected individuals to both pyogenic infections and rheumatic disorders. A defect in a complement regulatory protein, C1-inhibitor, results in the clinical disorder known as hereditary angioedema (14,15). A more complete discussion of identified deficiencies in complement components, and the disorders to which they relate, can be found in Chapter 29.

Kinins

The kinins are a group of serum proteases whose ultimate product is bradykinin, an agent known to induce smooth muscle contraction, vasoconstriction, and increased permeability of smaller blood vessels (16–20). The kinin cascade is initiated by a number of by-products of tissue damage—collagen, cartilage, basement membranes—as well as by endotoxin and inorganic materials, which serve to activate factor XII (or Hageman factor, better known as a participant in the clotting cascade, as described below). Factor XII mediates the cleavage of prekallikrein to kallikrein, which not only serves to activate more factor XII, but also cleaves the proenzyme kininogen to produce bradykinin. Factor XII represents a crucial intersection, as it can also be activated by plasmin, a product of proteolytic cleavage among the fibrinolytic proteins, and by kallikrein, another protein of the kinin group.

Clotting and Fibrinolytic Proteins

In addition to the roles played by these proteins in hemostasis, they contribute significantly to the amplification of the inflammatory response via the direct activation of factor XII, as described above. Proteolytic cleavages initiated by activated factor XII ultimately result in the cleavage of fibrinogen to fibrin and to smaller fibrinopeptides, which serve as inflammatory modulators. Activated factor XII similarly activates the fibrinolytic system by generating the protease plasmin. Similar to factor XII, plasmin represents an important intersecting locus for all three protease systems, as its activity proceeds in a number of directions. Plasmin activity can generate fibrin split products, also inflammatory mediators, but more importantly, plasmin activity augments the production of activated factor XII, and in direct activation of the complement pathway via proteolytic cleavage of factor C3 (21–26).

Lipid Mediators

Lipid mediators are a complex group of chemicals that also participate in augmenting the inflammatory response. This group includes the prostaglandins, leukotrienes, platelet-activating factor (PAF), and a novel group known as lipoxins (27).

Prostaglandins

Prostaglandins are oxidized derivatives of the fatty acid arachidonate that mediate a number of the cardinal signs of inflammation, including fever, pain, and vascular permeability (28–33). The major sources of prostaglandin in acute inflammation include mononuclear phagocytes, endothelial cells, and platelets. Prostaglandin synthesis is augmented during inflammation by a number of stimuli, including bacterial endotoxin, immune complexes, complement component C3a, bradykinin, and IL-1. During inflammation, the production of prostaglandins is upregulated by a variety of mechanisms, including increased availability of fatty acid substrates, increased phospholipase activity, and increases in the level of cyclooxygenase activity. Prostaglandins mediate their proinflammatory effects through specific receptors present on target cells, and are known to promote the pain, fever, and edema characteristic of the acute inflammatory response.

Leukotrienes

Leukotrienes are also oxidation products of arachidonate that are synthesized in and released from neutrophils, and to a lesser extent, eosinophilic leukocytes. LTA₄ and its synthetic products LTB₄ and LTC₄ are synthesized and exported from these cells; LTA₄ can then be taken up by erythrocytes, platelets, and endothelial cells, where conversion to LTB₄ and LTC₄ can also take place. LTD₄ and LTE₄ are additional metabolic conversion products of LTC₄. Although evidence suggesting the existence for specific receptors for individual leukotrienes exists, these receptors remain undefined. Together, leukotrienes mediate a large array of proinflammatory activities, including vasoconstriction, increased vascular permeability, and increased endothelial adhesiveness, as well as neutrophil chemotaxis and activation (34–37). Most recently, leukotrienes have received attention as contributors to the pathophysiology of asthma (38–42).

Platelet-activating Factor

PAF (43–47) is a substituted derivative of glycerol phosphate that exists in both circulating and cellular forms. In its cellular form in endothelial cells, PAF has been shown to enhance neutrophil–endothelial cell adhesion. Specific receptors for PAF have been identified on neutrophils (48), and numerous antagonists have been identified (49–53).

Peptides and Amines

Histamine and Serotonin

Histamine, a decarboxylated derivative of the amino acid histidine was among the earliest of the soluble inflammatory mediators to be discovered (54–57). Tissue mast cells and basophils synthesize and store histamine, which is released in response to variety of physical and chemical stimuli (55). Histamine diffuses rapidly through tissues and into the bloodstream, and promotes many of the sequelae of acute inflammation, including vasodilation, increased vascular permeability, and interactions with the peripheral nervous system. As is the case with most other inflammatory mediators, histamine is recognized by specific receptors, in this case three distinct receptors, H₁, H₂, and H₃ (58–66). Serotonin, a derivative of tryptophan, is stored in platelets, mast cells, and ente-

rochromaffin cells of the gastrointestinal tract and is released through degranulation. Similar to histamine, serotonin has receptor-mediated vasoactive properties, although its role in acute inflammation is not well defined (54,67).

Neuropeptides

Neuropeptides are among the many components connecting the nervous system and the inflammatory response. As a group, neuropeptides are inflammatory mediators released from neurons in response to local tissue damage. This group of mediators includes substance P, vasoactive intestinal peptide, somatostatin, and calcitonin gene-related peptide (68–73). While numerous immunomodulatory activities have attributed to these mediators, the determination of the true physiologic roles and overall effects produced by these proteins is currently under study. Also under consideration are the roles played by neuropeptide-degrading enzymes, such as neutral endopeptidase, shown to be expressed on the neutrophil cell surface (74–76).

Nitric Oxide

Although its activity as both a neurotransmitter and an agent maintaining hemodynamic stability has been established, the role of nitric oxide in human host defense has been quite controversial (77–82). In initial studies, stimulation of human macrophages with lipopolysaccharide, interferon- γ (IFN- γ), granulocyte-macrophage colony-stimulating factor (GM-CSF), TNF- α , or heat-killed bacteria failed to elicit production of nitric oxide (83), in contrast to results obtained with a murine system. In contrast, a more recent study (84) demonstrated the generation of nitrite in human macrophage cultures in response to TNF- α and GM-CSF together with avirulent mycobacterial strains. Other groups have also demonstrated high levels of nitric oxide synthesis in response to a select group of stimuli (85–87). The molecular basis for this selectivity is currently an area of intense investigation.

Proinflammatory Cytokines

The identification and characterization of these soluble mediators has been one of the most active fields in current inflammation research (88–90). New cytokines are being discovered, and new activities for known cytokines are still emerging. It is often difficult to discuss the physiologic actions of an individual cytokine, as the interactions among cytokines and their cellular targets are complex. Rather than an exhaustive list, this is a brief overview of those mediators with major roles in the inflammatory response, and a review of some recent research focusing on the interplay of these mediators. A more comprehensive discussion of individual proinflammatory cytokines can be found in Chapter 22 and within the references listed.

Interleukin-1

IL-1 is a major inflammatory mediator, produced primarily by monocytes and activated macrophages (91–99). IL-1 activity is produced by two polypeptides (IL-1 α and IL-1 β) encoded by two distinct genes on chromosome 2. High-affinity receptors for IL-1 are found on lymphocytes and fibroblasts. Numerous local and sys-

temic proinflammatory activities have been attributed to IL-1, including increasing local blood flow, fever, production of other soluble mediators, and enhanced expression of adhesion molecules. An unusual feature of IL-1 is the presence of a naturally occurring antagonist, IL1RA, which is expressed in neutrophils and monocytes (100,101).

Interleukin-4

IL-4 has a number of activities related to allergic inflammation, including stimulating basophil development, eosinophil chemotaxis, and expression of IgE receptors on B cells (102–108). IL-4 also participates in cell fusion related to the formation of granulomas, and also has antiinflammatory properties.

Interleukin-6

IL-6 is produced by T-lymphocytes, endothelial cells, monocytes, and fibroblasts, and has wide-reaching effects on T- and B-lymphocytes and macrophages, including promoting monocyte differentiation, increased number of circulating platelets, and synthesis of acute phase reactant proteins (including fibrinogen) in the liver (102,109–113).

Interleukin-8

IL-8 is a chemokine whose synthesis can be induced in a variety of cell types (monocytes, lymphocytes, and neutrophils) stimulated with IL-1 α , IL-1 β , or TNF. The activities of IL-8, however, appear to be restricted to neutrophils, enhancing both the chemotactic and degranulation responses. At the molecular level, IL-8 induces a net increase in the expression of cell surface adhesion molecules and elicits activation of the neutrophil NADPH oxidase (114–121).

Tumor Necrosis Factor

TNF (122–128) is derived from activated macrophages; TNF- α and TNF- β are two distinct but related polypeptides. TNF is associated with the production of fever and, similar to IL-1, promotes the increased expression of most other proinflammatory mediators, and it is prominently associated with the induction of cellular apoptosis (129–131).

Interferon- γ

IFN- γ (132–137) is a product of T cells and natural killer (NK) cells. Although initially recognized as an antiviral agent, IFN- γ activities are wide-ranging; best characterized is its ability to increase generation of highly reactive oxygen species such as superoxide anion and hydrogen peroxide and to alter the cell surface antigens of macrophages, permitting them to eliminate invading pathogens. IFN- γ also mediates activities of endothelial cells and, to a somewhat lesser extent, neutrophils. IFN- γ has been shown to have utility in preventing infections in certain patients with compromised host defenses or infections with intracellular pathogens (135). Research on a family with enhanced susceptibility to mycobacterial infection has demonstrated a role for the IFN- γ receptor in the pathogenesis of this disease (138,139).

Interleukin-12

IL-12 is a heterodimeric product of macrophages and B-lymphocytes that enhances the synthesis of IFN- γ and stimulates proliferation of NK, T helper 1 (Th1) cells, and cytotoxic T-lymphocytes. The role of IL-12 in host defense against intracellular bacteria is discussed in Chapter 40.

MODEL SYSTEMS OF INFLAMMATION

Two model systems have provided insight into the temporal appearance and importance of mediators in the various processes of inflammation in humans. In one model, small amounts of the lipid-A derivative of endotoxin are administered intravenously to normal human subjects, and mediator accumulation in peripheral blood is monitored (140). In the other model, mediator accumulation is monitored locally in the skin following creation of skin blisters by suction (141,142).

Response to Intravenous Endotoxin

Following intravenous endotoxin, a characteristic change in body temperature and white blood count is observed. Body temperature begins to increase after about 1 hour and reaches a maximum at about 4 hours. The leukocyte count shows a characteristic decrease at about 30 minutes, due to neutrophil and monocyte adherence to endothelial cells in the lung and spleen. This is followed by a leukocytosis characterized by the presence of immature neutrophils at about 4 hours, which can persist throughout 24 hours, with gradual return to baseline by 48 hours. The leukocytosis is predominantly due to mobilization of immature neutrophils from the bone marrow. The critical components of the inflammatory response—fever, neutrophil margination in the circulatory vessels, and then mobilization from the bone marrow—are associated with readily detected changes in circulating levels of certain mediators of inflammation. For example, TNF- α peaks within 2 hours (143) and is likely the predominant pyrogen associated with the febrile response. Plasma levels of the chemoattractant IL-8 increase early and peak by 4 hours. Early increases in IL-8 may relate to the transient decrease in the neutrophil count at 30 minutes (margination), since administration of intravenous chemoattractants in experimental animals is associated with a rapid neutropenia, likely a result of increased neutrophil expression of adhesion receptors (CR3) (144). In this regard, it is of interest that significant increases in plasma C5a and LTB₄ were not detected following administration of intravenous endotoxin to humans, reinforcing the probable critical role of IL-8 in the process of neutrophil margination. Plasma concentrations of IL-6 also increased 2 to 4 hours following intravenous endotoxin. In addition to not detecting increases in plasma C5a, LTB₄, or IL-1, no increases in circulating IL-2, IL-3, IL-4, IFN- γ , transforming growth factor- β (TGF- β), or nitrate-nitrite were detected following intravenous endotoxin, emphasizing the specificity of the responses observed. Kuhns and colleagues (145) have described a patient with recurrent bacterial infections that displayed hyporesponsiveness to both endotoxin and IL-1, attributable to a defect in signal transduction. Although the precise molecular basis of endotoxin activity has not been determined, there is evidence to suggest that endotoxin interacts with the cell surface antigen CD14 on the surface of phagocytes (146,147) and modulates the expression of nuclear factor κ B (NF- κ B), a factor that promotes the transcription of numerous proinflammatory mediators (148).

Temporal Analysis of Soluble Mediators in Blister Fluid

Soluble mediator accumulation at local inflammatory processes can be detected in raised skin blisters induced in normal volunteers (141). Mediators detected in blister fluid within 3 to 5 hours of the inflammatory response included LTB₄, C5a, IL-8, and IL-6. In contrast, IL-1 β , GM-CSF, and TNF- α were not detected until after 8 hours in the blister. Although IFN- γ was reported by Kuhns and colleagues (141) to be an early mediator in skin blister fluid, these results have not been repeated in subsequent studies (D.B. Kuhns and J.I. Gallin, unpublished results). Small amounts of IL-4 accumulated into the skin blisters, but IL-2 and IL-1 α were not detected. Thus, the endotoxin and skin blister models of inflammation demonstrate that there are clear differences in the mediators that can be detected systemically and locally.

CELLULAR MEDIATORS OF THE ACUTE INFLAMMATORY RESPONSE

There are many who would argue that the entire interconnecting network of proinflammatory responses are designed to facilitate the recruitment of neutrophils to the site of tissue injury. Neutrophils are among the cells known as "professional phagocytes"; they respond to the soluble inflammatory mediators by migrating to the site of tissue injury and by ingesting and destroying invading pathogens and damaged tissue, leading the way to resolution and, ultimately, tissue repair. The recruitment of leukocytes (neutrophils, monocytes, and eosinophils) to foci of inflammation can be monitored by a both blister fluid and Rebuck skin window techniques, as discussed above. Other participants in the acute response include platelets, lymphocytes, and endothelial cells.

Neutrophils

Neutrophilic leukocytes are crucial to both immunity and inflammation, and prolonged neutropenia leads to inevitable demise as a result of overwhelming infection (149–151). Neutrophils normally represent between 40% and 50% of the circulating leukocyte population, and they are easily recognized on a Wright's stained blood smear by their size, their characteristic multilobed nuclei, and the presence of fine stippling, representing granules throughout the cytoplasmic compartment (Fig. 2). Primary and secondary granules contain distinct sets of their own proinflammatory mediators, as described below.

Development in the Bone Marrow

Neutrophils develop from undifferentiated precursors present in the bone marrow (152). The myeloblast is the first morphologically identifiable precursor of the neutrophil lineage, followed by the promyelocyte, myelocyte, metamyelocyte, and band form, which directly precedes the mature neutrophil (Fig. 3). Several products of activated T-lymphocytes, including GM-CSF, IL-3, and granulocyte colony-stimulating factor (G-CSF), participate in the process of neutrophil maturation by direct interactions with their respective receptors, which have been identified on neutrophil precursor cells. When mature, neutrophils are released into the circulation, where they can respond to the soluble proinflammatory mediators described above. Neutropenia may be related to chemotherapeutic




FIG. 2. A: Light and (B) electron microscopic views of a mature human peripheral blood neutrophil. (Illustrations courtesy of Dr. Douglas Kuhns.)

agents, autoantibodies, or infection. These conditions are often reversible; inherited disorders known as cyclic neutropenia and familial neutropenia have also been identified (153–157).

Activation and Priming

Neutrophils in the circulation are quiescent cells with only the potential to mediate a wide range of inflammatory activities. This potential is realized when neutrophils are activated (158,159). Neutrophils can be activated by a large (and increasing) number of specific agents (Table 1). As a group, these activating agents transmit signals to the neutrophils via interaction with specific cell surface receptors (149,158–164), many of which interact with intracellular components known as G proteins. G proteins catalyze the hydrolysis of guanosine triphosphate (GTP) to guanosine diphosphate (GDP) and inorganic phosphate, meanwhile initiating a series of events including activation of phospholipase C, initiation of calcium fluxes, and membrane depolarization. Once activated, neutrophils are able to adhere to endothelial cells, migrate through the endothelial barrier, and ingest and at least attempt to destroy

pathogens, foreign bodies, and remnants of tissue damage. An intriguing aspect of neutrophil activation is the phenomenon of priming. Neutrophils primed by brief exposure to activating agents (endotoxin, IL-1, f-Met-Leu-Phe, GM-CSF) exhibit an enhanced response to subsequent stimuli. Both short-term (including changes in cell shape, oxidative and phagocytic capacity) and long-term (prolonged cell viability) responses to priming agents have been observed. Overall, the phenomenon of priming suggests that neutrophil activation is a two-step process, requiring an initial switch from a nonreceptive to a receptive state. The molecular basis for this switch is currently under investigation.

Adherence

To participate effectively in the inflammatory process, neutrophils must ultimately leave the bloodstream and migrate into the tissues. The initial step in this process is adherence to the vascular endothelium. Neutrophil adherence is a two-step process, the first involving a class of cell surface molecules known as selectins (165–173). Selectins mediate the process in which neutrophils roll




FIG. 3. Stages of neutrophil maturation in the bone marrow, from myeloblast to mature neutrophil. The large black circles represent the primary granules, and the dark shading, the secondary granules. The stages at which the proinflammatory mediators myeloperoxidase and lactoferrin are synthesized are as indicated by the arrows. (Adapted from ref. 300.)

TABLE 1. Agents promoting neutrophil activation

Agent	Function stimulated
LTB ₄	Chemoattractant; enhances adherence to endothelial cells; activates degranulation and NADPH oxidase activity
Complement fragment C5a	Chemoattractant; induces degranulation and adherence
PAF	Induces aggregation and adherence, chemoattractant degranulation
Histamine	Concentration-dependent changes in chemotaxis priming, and degranulation
IFN- γ	Increases antibody-dependent cytotoxicity, priming
G-CSF	Increases antibody-dependent cytotoxicity, priming; enhances phagocytosis; stimulates maturation within bone marrow
GM-CSF	Priming; stimulates maturation within bone marrow
TNF- α	Chemoattractant, priming; enhances phagocytosis and antibody-dependent cytotoxicity
IL-8	Chemoattractant; induces degranulation and NADPH oxidase activity
fMet-Leu-Phe	Chemoattractant; induces aggregation, degranulation, and NADPH oxidase activity

or slow down prior to their actual activation-dependent adherence to the endothelial cells. There are three classes of selectins that have been identified: L-selectins, which have been identified on all leukocytes; E-selectin, on the surface of activated endothelial cells; and P-selectin, found on endothelial cells and platelets. Selectins function by binding to carbohydrate ligands present on the adhering cell. The ligand for the endothelial E-selectin is sialylated Lewis-X antigen, found on the neutrophil, which, when absent, results in a marked immunodeficiency state.

The second part of the adherence process is the tight binding mediated by integrins (174–180). The leukocyte integrins are a subgroup of an extensive family of proteins that mediate a wide range of interactions between cells, and between cells and the extracellular environment. The leukocyte integrins—LFA-1 (CD11a/CD18), Mac-1 (CD11b/CD18), and p150,95 (CD11c/CD18)—are heterodimeric proteins with distinct α and shared β polypeptide chains. Mac-1 in particular has several well-characterized roles in the inflammatory process. Mac-1 is stored in the secondary granule compartment and is brought to the cell surface in conjunction with neutrophil activation. In addition to mediating specific adherence, Mac-1 participates in phagocytosis and chemotaxis, and in production of reactive oxygen species (see below). Intercellular cell adhesion molecule-1 (181), a cell surface protein found on endothelial cells, has been identified as a ligand for Mac-1.

Two forms of inherited defects of neutrophil adhesion have been identified: Leukocyte adhesion deficiency I (LAD I) involves a genetic defect in the biosynthesis of CD18, the shared β chain for all three leukocyte adhesion molecules (182–184). The defect is autosomal recessive and has been mapped to human chromosome 22q22.3. Individuals with this disorder have frequent and recurrent skin and soft-tissue infections, poor wound healing, and severe periodontal disease. In contrast, leukocyte adhesion deficiency II (LAD II) is a glycosylation defect that results in the inability to synthesize the sialyl-Lewis X carbohydrate ligand for E- and P-selectin (185,186). This condition results in a defect in neutrophil rolling that occurs prior to and facilitates neutrophil adherence to endothelial cells, and affected individuals likewise suffer from frequent severe bacterial infections.

Chemotaxis

As part of the activation process, neutrophils are capable of sensing and responding to concentration gradients of the activating agents that are highlighted in Table 1. By “crawling” across a surface, neutrophils can be seen to migrate toward a higher concen-

tration of attractant. At the subcellular level, cell motility requires alterations in the neutrophil cytoskeleton, which is composed primarily of actin filaments. Although the precise mechanism by which signals are transmitted directly to the cytoskeleton is unclear, there is evidence to suggest that several actin-binding proteins (including profilin, cofilin, and gelsolin) participate in altering the actin filament structure, permitting net movement of the cell in response to a chemoattractant gradient (187–189).

Phagocytosis

Phagocytosis, or engulfment of foreign or damaged material, is the centerpiece of the inflammatory process and is discussed in detail in Chapter 30 (190,191). To engulf a particle, neutrophils extend pseudopodia, which encircle the offending material; the pseudopodia fuse, trapping the material inside the cell in a compartment known as a phagosome. Particles coated with immunoglobulins (or opsonized) are phagocytosed in a highly efficient fashion, as they are recognized by and bind directly to the Fc receptors present on the neutrophil cell surface. Particles opsonized by proteolytic products of complement (see above) are similarly phagocytosed in a specific, receptor-mediated fashion, involving CR1 and CR3.

Degranulation

The primary and secondary granules of neutrophils contain a number of distinct effector proteins, listed in Table 2. As part of the activation process, the cytoplasmic membrane-bound granules fuse with the phagosome, placing the effector proteins in direct contact with the ingested material. Among the highlights of the components of the primary (also known as azurophil) granules are lysozyme, which can digest the peptidoglycan component of most bacterial cell walls, and cathepsin G, defensins, and bacterial permeability-increasing protein (BPI), all with inherent antibacterial activity. Goldman and colleagues (192) have shown that human β -defensin-1 is inactivated in individuals with cystic fibrosis, and thus may be related to the pathogenesis of bacterial infections in affected individuals. Also among the more prominent components is myeloperoxidase, which converts hydrogen peroxide generated by the NADPH oxidase (see below) and hydrochloric acid to hypochlorous acid, another antimicrobial agent.

The secondary (or specific) granules contain several proteins whose role in the inflammatory response remains a bit mysterious.

TABLE 2. Major components of neutrophil primary and secondary granules

Primary granules	Secondary granules
Myeloperoxidase	Lactoferrin
Defensins	Gelatinase
BPI	Collagenase
Cathepsin G	Vitamin B 12-binding protein
Lysozyme	Lysozyme
Elastase	Cytochrome b558
Alkaline phosphatase	fMLP receptor
Proteinase 3	CD11b/CD18, CD11c/CD18 (integrins)
Beta glucuronidase	Complement receptor 3 (CR3)
Alpha fucosidase	Histaminase
Phospholipases A2, C, D	Plasminogen activator
Alpha mannosidase	

Among these proteins is lactoferrin, an iron-binding protein with some antibacterial activity (193). The secondary granules also contain stored sources of CR3 and other receptors for neutrophil activation agents, as well as stored membrane components of the NADPH oxidase.

Chediak-Higashi syndrome is a disorder in which neutrophils demonstrate abnormal morphology, abnormal chemotaxis, and failure to degranulate, and affected individuals are subjected to recurrent, severe bacterial and fungal infections. Three independent groups have reported the identification of the genetic defect in Chediak-Higashi syndrome, residing on human chromosome 1q42-43 (194-196). Disorders of neutrophils are specific granule deficiency, in which the secondary, or specific, granules are absent or, alternatively, are present, but without the granule protein components (197-199) and myeloperoxidase deficiency (200-202).

NADPH Oxidase

A crucial component of the neutrophil host defense mechanism is the enzyme complex known as the NADPH oxidase (Fig. 4) (203-210). This enzyme assembles on the phagosomal membrane

from two integral membrane components (gp91-phox and p22-phox. for phagocyte oxidase) and three cytosolic components (p47-phox, p67-phox, and rac) to catalyze the production of superoxide anion from molecular oxygen and free electrons; superoxide is then converted to the toxic metabolite hydrogen peroxide by the actions of superoxide dismutase, or to hypochlorous acid by the primary granule protein myeloperoxidase.

While the ability to generate toxic oxygen metabolites is crucial to host defense, these agents represent the sharpest part of the double-edged sword of inflammation. Superoxide anion is readily diffusible through membranes and can be converted to toxic metabolites outside the restricted locale of the phagosome. Products of oxygen radicals can create enlarged foci of tissue damage, thereby enhancing and augmenting the inflammatory process far beyond what was necessary to contain the initial insult. Similarly, oxidative injury has been implicated in the pathogenesis of cardiovascular, neoplastic, arthritic, and neurodegenerative disease. Chronic granulomatous disease is an inherited disorder in which neutrophils are rendered incapable of generating toxic oxygen metabolites (207,211-213). This results in an inability to mount an effective defense against bacteria (particularly catalase-positive strains) and fungi, and affected individuals often present with recurrent, life-threatening infections. Inherited defects in any one of the four oxidase proteins can disable the enzyme complex and result in disease. Therapy for this disorder includes prophylactic antibiotics and injections of the inflammatory modulating agent IFN- γ (214,215).

Protein Biosynthesis

Although neutrophils are generally perceived as "end-stage" cells, more recent work has shown that neutrophils are indeed capable of significant biosynthetic activity. The proteins expressed *de novo* in neutrophils include components of the NADPH oxidase and specific membrane receptors and antigens. Several proinflammatory mediators are released by activated neutrophils, including IL-1, TNF- α , IL-6, IL-8, GM-CSF, G-CSF, and plasminogen activator (149,211-215). Kuhns and Gallin (216) have shown that IL-8 is actively synthesized by exudate neutrophils. These mediators can "feed back" on the system and augment the overall inflammatory response.




FIG. 4. Schematic of the protein components of the phagocyte NADPH oxidase. On activation, the cytoplasmic components rac, p67, p47, and p40 are translocated to the cell membrane to form the catalytic complex. Once formed, the complex can catalyze the conversion of molecular oxygen (O_2) to superoxide (O_2^-), which is then converted to the toxic oxygen metabolite, hydrogen peroxide (H_2O_2) by the actions of superoxide dismutase (SOD). The proteins associate via specific recognition sites, known as SH3 domains, as described in refs. 301-303. (Illustration courtesy of Dr. Thomas Leto.)

Monocytes and Macrophages

Monocytes, also within the "professional phagocyte" grouping, migrate into the tissues, as do the neutrophils (217–223). Activated macrophages, like neutrophils, are capable of phagocytosis and release antibacterial proteins and proinflammatory mediators. Macrophage functions complement those of neutrophils during the acute response, and they take on a more central role during chronic inflammation. The biology and physiology of macrophages are covered in detail in Chapter 15.

Eosinophils

Eosinophils are primarily tissue-dwelling granulocytes that are also recruited to sites of acute inflammation, seen most prominently in response to respiratory, gastrointestinal, and dermatologic allergens, and to generalized infection with helminthic parasites (224–228). Similar to neutrophils, eosinophils develop in the bone marrow, have receptor-mediated responses to specific activating agents (including RANTES, MIP-1 α , and eotaxin), and contain cytoplasmic granules with oxidative and cationic proteins; in contrast to neutrophils, eosinophils are ineffective phagocytes and release the contents of their granules to the extracellular milieu. Interestingly, the detrimental features of eosinophils are among the best characterized, particularly their contributions to the pathogenesis of reactive airway disease (229–231), while the beneficial aspects of eosinophils in the inflammatory response remain poorly (if at all) understood.

Platelets and Lymphocytes

Platelets contribute to the inflammatory response by several different mechanisms (236–247). Platelets contain and can release numerous inflammatory mediators, including fibrinogen, plasminogen, and other components that participate in the plasma protease systems, lipids, and serotonin. Several of these mediators released from platelets are direct activating agents for neutrophils, and, conversely, mediators released by activated neutrophils (oxygen metabolites, granule proteins, lipids) serve to alter platelet function. Platelets interact with lymphocytes, providing the cell contact and reagents for prostaglandin synthesis. Platelets also interact with fibroblasts, stimulating collagen and fibronectin synthesis during resolution (see later discussion).

The complex biology of T- and B-lymphocytes and their role in specific immunity are discussed in detail in Chapters 6, 7, 10 and 11; the role of immunoglobulins in augmenting the inflammatory response and enhancing neutrophil phagocytosis has been discussed above. It is also important to emphasize that many of the soluble mediators discussed in the earlier sections (interleukins, IFN- γ) are produced by activated T-lymphocytes, as are a number of the antiinflammatory mediators to be discussed below.

Endothelial Cells

The role of endothelial cells in providing a base for neutrophil adherence has already been discussed. More recently appreciated is the fact that endothelial cells synthesize and release numerous proinflammatory mediators (236–247; see also references from selectins and integrins above).

ALLERGY AND INFLAMMATION

Allergy, also known as the immediate hypersensitivity response, is also a form of inflammation and is considered in detail in Chapter 35. The central components of this type of the allergic response are IgE, IgE receptors on basophils and mast cells, and histamine released from these cells upon IgE receptor-mediated interaction. The role of allergy in host defense remains controversial and is focused on the role of IgE and mucosal mast cells in the defense against gastrointestinal parasites (248–250); most of the literature on allergy focuses on its detrimental features.

RESOLUTION OF THE ACUTE INFLAMMATORY RESPONSE

Resolution, or the way in which the acute inflammatory response is downregulated, is currently an area of active research. The mediators promoting inflammatory resolution may ultimately be harnessed for use as therapeutic agents in limiting the injurious aspects of acute inflammation.

Cell Senescence or Apoptosis

A concept that has been appreciated only recently, apoptosis, or programmed cell death, is an active process in which cells, responding to specific stimuli, undergo a stereotypical pattern of morphologic changes (nuclear condensation, DNA "laddering") prior to their eventual demise. Granulocyte apoptosis as a means of inflammatory resolution is an intriguing avenue of current research (251–257). Several cytokines have been reported to modulate neutrophil apoptosis *in vitro*, including TNF- α , eicosanoids, IL-10, and antioxidants (258–262); the role of these mediators in the resolution of acute inflammation awaits future study (see also Chapter 23).

Antiinflammatory Mediators

The antiinflammatory effects of a number of soluble mediators have been characterized. There are most likely many others awaiting identification and characterization (see also Chapter 22).

Interleukin-4

In addition to the proinflammatory effects described previously, IL-4 downregulates IL-6 production and is involved in the downregulation of neutrophil superoxide production (263).

Transforming Growth Factor- β

TGF- β promotes several antiinflammatory effects, including suppression of hematopoiesis, reduction in production of proinflammatory cytokines, and inhibition of leukocyte adhesion (264–269). Perhaps most dramatic, TGF- β 1 knock-out mice develop severe inflammation in multiple tissues, suggesting the primary role of TGF- β as that of an antiinflammatory mediator (270,271). TGF- β is produced in many cell types, including platelets, macrophages, and T- and B-lymphocytes.

Interleukin-10 and Interleukin-13

IL-10 is produced by macrophages and CD8⁺ T-lymphocytes, and it has been shown to inhibit the activation of specific macrophage

subsets, including inhibition of the production of proinflammatory cytokines and interference with the macrophage-mediated antigen presentation. IL-10 is also implicated in host response to both Epstein-Barr virus and human immunodeficiency virus infection (272–276). IL-13 has been observed to induce IL-4-independent IgE synthesis and to induce proliferation and differentiation of human B cells activated by the CD40 ligand (277–279).

Hypothalamo–Pituitary–Adrenocortical Axis

One of the more intriguing avenues of investigation is the connection between the central nervous system, the adrenal cortex, and the resolution of inflammation (280–284). An appreciation of this phenomenon relates to the observation that glucocorticoids, produced by the adrenal cortex, mediate immunosuppression, and thus may downregulate the acute inflammatory response. Numerous studies have suggested that IL-1, IL-6, and TNF- α promote marked increases in hypothalamic stimulation, leading to increases in serum ACTH and corticosterone in experimental animal systems; prostaglandins have also been implicated in this process.

Wound Repair and Angiogenesis

Several morphologic stages of wound repair have been described (264–268). Neutrophils and macrophages carry out the initial debridement, including removal of foreign material and cellular debris. Fibroblasts and epithelial and endothelial cells, responding to multiple inflammatory mediators, grow and divide to create new tissue and restore function. Angiogenesis is the process by which new tissue is revascularized. The formation of capillaries has been shown to proceed through several well-defined morphologic events, including vasodilation of the parent venule or capillary, removal of the preexisting basement membrane, migration and proliferation of endothelial cells, and formation of a new lumen. These events are promoted by numerous soluble mediators, including epidermal growth factor, keratinocyte growth factor, platelet-derived growth factor, fibroblast growth factors, TGF- α , TGF- β , and cellular mediators (macrophages, platelets, keratinocytes, endothelial cells, and mast cells) (285–289).

CHRONIC INFLAMMATION

When acute inflammation persists, either through incomplete clearance of the initial inflammatory focus or as a result of multiple acute events occurring in the same location, chronic inflammation takes over. In contrast to acute inflammation, which is characterized by a primarily neutrophil influx, the histologic findings in chronic inflammation include accumulation of macrophages and lymphocytes and growth of fibroblasts and vascular tissue. It is these latter two features that result in the tissue scarring that is typically seen at sites of prolonged or repeated inflammatory activity.

Among the most interesting sequelae of chronic inflammation is the formation of a tissue granuloma (290–296). A granuloma is a collection inflammatory cells—principally macrophages and lymphocytes, which are eventually surrounded by a fibrotic wall—that forms in tissues as part of the inflammatory response to a persistent irritant. Several unusual cell types are characteristic of granulomata, including epithelioid cells, which are macrophage derivatives, and multinuclear giant cells, which are fusions of epithelioid cells and




FIG. 5. Light microscopic image of a tissue granuloma from the murine CGD (p47 phox knock-out) model (304). Lymphocytes comprise the central core, which is surrounded by macrophages and fibroblasts.

macrophages (Fig. 5). Although the precise mechanism of granuloma formation and resolution is not yet clear, the actions of specifically sensitized T cells and their soluble mediators (including TNF- α and IFN- γ) participate in the formation and maintenance of granulomata in their active state. Several conditions predispose an individual to granuloma formation, most notably the presence of intracellular bacteria (e.g., tuberculosis; see Chapter 40) or inorganic antigens (e.g., berylliosis).

More recent work on the molecular mechanisms underlying chronic inflammation has focused on the role of NF- κ B, a transcription factor originally identified as regulating the expression of the murine κ light chains. Since that time, NF- κ B activation has been associated with endotoxin, cytokines, viruses, and oxidants, and NF- κ B has been shown to regulate expression of adhesion molecules, E-selectin, and numerous chemotactic cytokines (297–299).

FUTURE DIRECTIONS: NOVEL ANTIINFLAMMATORY THERAPIES

The goal of antiinflammatory therapy is to eliminate the undesirable aspects of the double-edged sword—tissue destruction beyond what is absolutely necessary for containing and eliminating a pathogenic agent. At the same time, antiinflammatory therapy must be sufficiently short-lived and/or selective so as to avoid creating an immunocompromised host. To this end, several generalized antiinflammatory agents (e.g., glucocorticoids, nonsteroidal antiinflammatory agents) have been recognized for their broad scope of effectiveness. Specific agents on the horizon may be more effective at pinpointing specific aspects of the inflammatory response that might be more carefully controlled.

Immunophilin-binding Agents

Cyclosporine (CsA) and FK506 (tacrolimus) are the most current of this group of generalized antiinflammatory agents (305–310). CsA and FK506 are structurally unrelated agents that selectively inhibit T-lymphocyte activation by interfering with the transcription of several cytokine genes. In the most recently proposed molecular mechanism, CsA and FK506 interact with distinct intracellular-binding proteins, known as immunophilins, which have enzymatic activity that appears to be unrelated to their role in immunosuppression. The immunophilin-bound complexes formed interfere with calcineurin-mediated transcriptional activation, thus inhibiting transcription of the genes encoding the proinflammatory agents IL-2, IL-3, IL-4, GM-CSF, IL-8, and IFN- γ ; the complexes also function by suppressing T-cell proliferation. Although used primarily as adjuvant therapy in organ transplantation, they have also been used in the treatment of rheumatoid arthritis and inflammatory bowel disease (311–314).

Experimental Antiinflammatory Agents

Several groups have experimented with agents that specifically inhibit individual proinflammatory mediators (315–332) or pathways (333–337), including specific cytokine and receptor antagonists. Other approaches take their lead from the actions of the immunophilin-binding proteins, and have considered ways to exert transcriptional control on one or more proinflammatory mediators (337,338). In the not too distant future, gene replacement therapy may emerge as an option, once the genetic bases of the complex inflammatory disorders have been identified and elucidated.

REFERENCES

- Hunter J. *A treatise of the blood inflammation, and gunshot wounds*. vol I. London: J. Nicoll. 1794.
- Cohnheim J. *Lectures in general pathology*. vol I. London: New Sydenham Society, 1889. McKee AD. translator, from the German edition.
- Dale HH, Laidlaw PP. The physiologic action of b-imidazolylethylamine. *J Physiol* 1911;41:318–344.
- Gallin JI, Goldstein IM, Snyderman R, eds. *Inflammation*. 2nd ed. New York: Raven Press, 1992.
- Musher D, Cohen M, Baker C. Immune responses to extracellular bacteria. In: Rich RR, Fleisher TA, Schwartz BD, Shearer WT, Strober W, eds. *Clinical immunology*. vol I. St. Louis: Mosby-Year Book, 1996:479–502.
- Robbins SL, Cotran RS. Inflammation and repair. In: *Pathologic basis of disease*. 2nd ed. Philadelphia: WB Saunders, 1979:59–63.
- Ward PA. Role of complement, chemokines, and regulatory cytokines in acute lung injury. *Ann NY Acad Sci* 1996;796:104–112.
- Goldstein IM. Complement: biologically active products. In: Gallin JI, Goldstein IM, Snyderman R, eds. *Inflammation*. 2nd ed. New York: Raven Press, 1992: 63–80.
- Kozel TR. Activation of the complement system by pathogenic fungi. *Clin Microbiol Rev* 1996;9:34–46.
- Muller-Eberhard HJ. Complement: chemistry and pathways. In: Gallin JI, Goldstein IM, Snyderman R, eds. *Inflammation*. 2nd ed. New York: Raven Press, 1992:33–62.
- Holers VM. Complement. In: Rich RR, Fleisher TA, Schwartz BD, Shearer WT, Strober W, eds. *Clinical immunology: principles and practice*. St. Louis: Mosby-Year Book, 1996:363–391.
- Holers VM. Complement. In: Rich RR, Fleisher TA, Schwartz BD, Shearer WT, Strober W, eds. *Clinical immunology: principles and practice*. St. Louis: Mosby-Year Book, 1996:363–391.
- Ratnoff WD. Inherited deficiencies of complement in rheumatic diseases. *Rheum Dis Clin North Am* 1996;22:75–94.
- Ono H, Kawaguchi H, Ishii N, Nakajima H. A point mutation in exon 7 of the C1-inhibitor gene causing type I hereditary angioedema. *Hum Genet* 1996;98: 452–453.
- Ernst SC, Circolo A, Davis AE, Gheesling-Mullis K, Fliesler M, Strunk RC. Impaired production of both normal and mutant C1 inhibitor proteins in type I hereditary angioedema with a duplication in exon 8. *J Immunol* 1996;157: 405–410.
- Busse R, Fleming I. Molecular responses of endothelial tissue to kinins. *Diabetes* 1996;45[Suppl 1]:S8–S13.
- Gepetti P, Bertrand C, Ricciardolo FL, Nadel JA, Ricciardolo FL. New aspects on the role of kinins in neurogenic inflammation. *Can J Physiol Pharmacol* 1995;73:843–847.
- Margolius HS. Kallikreins and kinins. Molecular characteristics and cellular and tissue responses. *Diabetes* 1996;45[Suppl 1]:S14–S19.
- Margolius HS. Kallikreins and kinins. Some unanswered questions about system characteristics and roles in human disease. *Hypertension* 1995;26:221–229.
- Mombouli JV, Vanhoutte PM. Kinins and endothelial control of vascular smooth muscle. *Annu Rev Pharmacol Toxicol* 1995;35:679–705.
- Carmeliet P, Collen D. Gene targeting and gene transfer studies of the biological role of the plasminogen/plasmin system. *Thromb Haemostasis* 1995;74:429–436.
- Altieri DC. Inflammatory cell participation in coagulation. *Semin Cell Biol* 1995;6:269–274.
- Esmon CT, Fukudome K. Cellular regulation of the protein C pathway. *Semin Cell Biol* 1995;6:259–268.
- Esmon CT. Inflammation and thrombosis: the impact of inflammation on the protein C anticoagulant pathway. *Haematology* 1995;80[Suppl 2]:49–56.
- Salgado A, Boveda JL, Monasterio J, et al. Inflammatory mediators and their influence on haemostasis. *Haemostasis* 1994;24:132–138.
- Benedict CR, Pakala R, Willerson JT. Endothelial-dependent procoagulant and anticoagulant mechanisms. Recent advances in understanding. *Tex Heart Inst J* 1994;21:86–90.
- Brady HR, Serhan CN. Lipoxins: putative braking signals in host defense, inflammation and hypersensitivity. *Curr Opin Nephrol Hypertens* 1996;5:20–27.
- Davies P, MacIntyre DE. Prostaglandins and inflammation. In: Gallin JI, Goldstein IM, Snyderman R, eds. *Inflammation*. 2nd ed. New York: Raven Press, 1992:123–138.
- Wallace JL, Tigley AW. New insights into prostaglandins and mucosal defense. *Aliment Pharmacol Ther* 1995;9:227–35.
- Vane JR, Botting RM. New insights into the mode of action of anti-inflammatory drugs. *Inflamm Res* 1995;44:1–10.
- Goetzl EJ, An S, Smith WL. Specificity of expression and effects of eicosanoid mediators in normal physiology and human diseases. *FASEB J* 1995;9:1051–1058.
- Seibert K, Masferrer J, Zhang Y, et al. Mediation of inflammation by cyclooxygenase-2. *Agents Actions Suppl* 1995;46:41–50.
- Asby B. Interactions among prostaglandin receptors. *Receptor* 1994;4:31–32.
- Lam BK, Austen KF. Leukotrienes: biosynthesis, release, and actions. In: Gallin JI, Goldstein IM, Snyderman R, eds. *Inflammation*. 2nd ed. New York: Raven Press, 1992:139–148.
- Ford-Hutchinson AW. Regulation of leukotriene biosynthesis. *Cancer Metastasis Rev* 1994;13:257–267.
- Henderson WR Jr. The role of leukotrienes in inflammation. *Ann Intern Med* 1994;121:684–697.
- Obata T, Yamashita N, Nakagawa T. Leukotriene and thromboxane antagonists. *Clin Rev Allergy* 1994;12:79–93.
- Busse WW. The role of leukotrienes in asthma and allergic rhinitis. *Clin Exp Allergy* 1996;26:868–879.
- Spector SL. Leukotriene inhibitors and antagonists in asthma. *Ann Allergy Asthma Immunol* 1995;75:463–470.
- Dahlen B, Dahlen SE. Leukotrienes as mediators of airway obstruction and inflammation in asthma. *Clin Exp Allergy* 1995;25:50–54.
- Harris RR, Carter GW, Bell RL, Moore JL, Brooks DW. Clinical activity of leukotriene inhibitors. *Int J Immunopharmacol* 1995;17:147–156.
- Israel E. Moderating the inflammation of asthma: inhibiting the production or action of products of the 5-lipoxygenase pathway. *Ann Allergy* 1994;72:279–284.
- Kingsnorth AN. Platelet-activating factor. *Scand J Gastroenterol Suppl* 1996: 219:28–31.
- Bussolino F, Camussi G. Platelet-activating factor produced by endothelial cells. A molecule with autocrine and paracrine properties. *Eur J Biochem* 1995;229: 327–337.
- Pinckard RN, Woodard DS, Showell HJ, Conklyn MJ, Novak MJ, McManus LM. Structural and (patho)physiological diversity of PAF. *Clin Rev Allergy* 1994;12: 329–359.
- Snyder F. Metabolic processing of PAF. *Clin Rev Allergy* 1994;12:309–327.
- Zimmerman GA, Prescott SM, McIntyre TM. Platelet-activating factor: a fluid-phase and cell-associated mediator of inflammation. In: Gallin JI, Goldstein IM, Snyderman R, eds. 2nd ed. *Inflammation*. New York: Raven Press, 1992:149–176.
- Dent G, Ukena D, Chanez P, Sybrecht G, Barnes P. Characterization of PAF receptors on human neutrophils using the specific antagonist, WEB 2086. *FEBS Lett* 1989 244:365–368.
- Hilger RA, Koller M, Konig W. Inhibition of leukotriene formation and IL-8 release by the PAF-receptor antagonist SM-12502. *Inflammation* 1996;20:57–70.
- Catalan RE, Martinez AM, Aragonés MD, Lombardía M, Garde E. PCA-4248, a PAF receptor antagonist, inhibits PAF-induced phosphoinositide turnover. *Eur J Pharmacol* 1995;290:183–188.
- Canz MJ, Weg VG, Walsh DT, Williams TJ, Nourshargh S. Differential effects of the PAF receptor antagonist UK-74, 505 on neutrophil and eosinophil accumulation in guinea-pig skin. *Br J Pharmacol* 1994;113:513–521.
- Yokota Y, Inamura N, Asano M, et al. Effect of FR128998, a novel PAF receptor antagonist on endotoxin-induced disseminated intravascular coagulation. *Eur J Pharmacol* 1994;258:239–246.

53. Underwood SL, Lewis SA, Raeburn D. RP 59227, a novel PAF receptor antagonist: effects in guinea pig models of airway hyperreactivity. *Eur J Pharmacol* 1992;210:97-102.
54. Atkinson TP, White MV, Kaliner MA. Histamine and serotonin. In: Gallin JI, Goldstein IM, Snyderman R, eds. *Inflammation*. 2nd ed. New York: Raven Press, 1992:193-210.
55. MacDonald SM. Histamine-releasing factors. *Curr Opin Immunol* 1996;8:778-783.
56. Raud J, Thorlacius J, Xie X, Lindbom L, Hedqvist P. Interactions between histamine and leukotrienes in the microcirculation. Aspects of relevance to acute allergic inflammation. *Ann NY Acad Sci* 1994;744:191-198.
57. Greaves MW, Sabroe RA. Histamine: the quintessential mediator. *J Dermatol* 1996;23:735-740.
58. Gothert M, Garbarg M, Hey JA, Schlicker E, Schwartz JC, Levi R. New aspects of the role of histamine in cardiovascular function: identification, characterization and potential pathophysiological importance of H3 receptors. *Can J Physiol Pharmacol* 1995;73:558-564.
59. Du Buske LM. Clinical comparison of histamine H1-receptor antagonist drugs. *J Allergy Clin Immunol* 1996;98:S307-S318.
60. Roberts DJ. A preclinical overview of ebastine. Studies on the pharmacological properties of a novel histamine H1 receptor antagonist. *Drugs* 1996;51:8-14.
61. Nielsen HJ. Histamine-2 receptor antagonists as immunomodulators: new therapeutic views? *Ann Med* 1996;28:107-113.
62. Leurs R, Vollinga RC, Timmerman H. The medicinal chemistry and therapeutic potentials of ligands of the histamine H3 receptor. *Prog Drug Res* 1995;45:107-165.
63. Leurs R, Smit MJ, Timmerman H. Molecular pharmacological aspects of histamine receptors. *Pharmacol Ther* 1995;66:413-463.
64. Zingel V, Leschke C, Schunack W. Developments in histamine H1-receptor agonists. *Prog Drug Res* 1995;44:49-85.
65. Arrang JM, Drutel G, Garbarh M, Ruat M, Traffort E, Schwartz JC. Molecular and functional diversity of histamine receptor subtypes. *Ann NY Acad Sci* 1995;757:314-323.
66. Arrang JM. Pharmacological properties of histamine receptor subtypes. *Cell Mol Biol* 1994;40:275-281.
67. Cazzola M, Matera MG, D Amato G, Rossi F. Effects of serotonin on airways: recent developments. *Allergy* 1995;50:1-10.
68. Chancellor-Freeland C, Zhu GF, Kage R, Beller DI, Leeman SE, Black PH. Substance P and stress-induced changes in macrophages. *Ann NY Acad Sci* 1995;771:472-484.
69. Hanesch U. Neuropeptides in dural fine sensory nerve endings—involvement in neurogenic inflammation? *Prog Brain Res* 1996;113:299-317.
70. Reubi JC, Laissue J, Waser B, Horisberger U, Schaer JC. Expression of somatostatin receptors in normal, inflamed and neoplastic human gastrointestinal tissues. *Ann NY Acad Sci* 1994;733:122-137.
71. Said SI. Vasoactive intestinal peptide and nitric oxide: divergent roles in relation to tissue injury. *Ann NY Acad Sci* 1996;805:379-387.
72. Weinstock JV. Vasoactive intestinal peptide regulation of granulomatous inflammation in murine Schistosomiasis mansoni. *Adv Neuroimmunol* 1996;6:95-105.
73. Reichlin S. Neuroendocrine-immune interactions. *N Engl J Med* 1993;329:1246-1253.
74. Matsumura T, Kugiyama K, Sugiyama S, et al. Neutral endopeptidase 24.11 in neutrophils modulates protective effects of natriuretic peptides against neutrophil-induced endothelial cytotoxicity. *J Clin Invest* 1996;97:2192-2203.
75. Painter RG, Aiken ML. Regulation of N-formyl-methionyl-leucyl-phenylalanine receptor recycling by surface membrane neutral endopeptidase-mediated degradation of ligand. *J Leukoc Biol* 1995;58:468-476.
76. Fagny C, Marchant A, De Prez E, Goldman M, Deschodt-Lanckman M. Lipopolysaccharide induced upregulation of neutral endopeptidase 24.11 on human neutrophils: involvement of the CD14 receptor. *Clin Sci* 1995;89:83-89.
77. Denis M. Human monocytes/macrophages: NO or no NO? *J Leukoc Biol* 1994;55:682-684.
78. Moncada S, Higgs A. The L-arginine-nitric oxide pathway. *N Engl J Med* 1993;329:2002-2012.
79. Schoedon G, Schneemann M, Walter R, Blau N, Hofer S, Schaffner A. Nitric oxide and infection: another view. *Clin Infect Dis* 1995;21[Suppl 2]:S152-S157.
80. Davies MG, Fulton GJ, Hagen PO. Clinical biology of nitric oxide. *Br J Surg* 1995;82:1598-1610.
81. Granger DL, Hibbs JB Jr. High-output nitric oxide: weapon against infection? *Trends Microbiol* 1996;4:46-47.
82. Lowenstein CJ, Dinerman JL, Snyder SH. Nitric oxide: a physiologic messenger. *Ann Intern Med* 1994;120:227-237.
83. Schneemann M, Schoedon G, Hofer S, Blau N, Guerrero L, Schaffner A. Nitric oxide synthase is not a constituent of the antimicrobial armature of human mononuclear phagocytes. *J Infect Dis* 1993;167:1358-1363.
84. Denis M. Tumor necrosis factor and granulocyte macrophage-colony stimulating factor stimulate human macrophages to restrict growth of virulent *Mycobacterium avium*: killing effector mechanism depends on the generation of reactive nitrogen intermediates. *J Leukoc Biol* 1991;49:380-382.
85. Hibbs JB, Westenfelder C, Taintor R, et al. Evidence for cytokine-inducible nitric oxide synthesis from L-arginine in patients receiving interleukin-2 therapy. *J Clin Invest* 1992;89:867-877.
86. Bukrinsky M, Schmidtmayerova H, Zybarth G, Dubrovsky L, Sherry B, Enikolopov G. A critical role of nitric oxide in human immunodeficiency type 1-induced hyperresponsiveness of cultures monocytes. *Mol Med* 1996;2:460-468.
87. Zembala M, Siedlar M, Marcinkiewicz J, Pryjma J. Human monocytes are stimulated for nitric oxide release in vitro by some tumor cells but not by cytokines and lipopolysaccharide. *Eur J Immunol* 1994;24:435-439.
88. Howard OM, Ben-Baruch A, Oppenheim JJ. Chemokines: progress toward identifying molecular targets for therapeutic agents. *Trends Biotechnol* 1996;14:46-51.
89. Murphy PM. Chemokine receptors: structure, function and role in microbial pathogenesis. *Cytokine Growth Factor Rev* 1996;7:47-64.
90. Prieschl EE, Kulmburg PA, Baumruker T. The nomenclature of chemokines. *Int Arch Allergy Immunol* 1995;107:475-483.
91. Estrov Z, Kurczok R, Talpaz M. Interleukin-1 and its inhibitors: implications for disease, biology and therapy. *Cancer Treat Res* 1995;80:51-82.
92. Dinarello CA. Interleukin-1 and interleukin-1 receptor antagonist. *Nutrition* 1995;11:492-494.
93. Dinarello CA. Biologic basis for interleukin-1 in disease. *Blood* 1996;87:2095-2147.
94. Fantuzzi G, Dinarello CA. The inflammatory response in interleukin-1 beta deficient mice: comparison with other cytokine-related knock-out mice. *J Leukoc Biol* 1996;59:489-493.
95. Saklatvala J. Intracellular signalling mechanisms of interleukin 1 and tumour necrosis factor: possible targets for therapy. *Br Med Bull* 1995;51:402-418.
96. Kupper TS, Groves RW. The interleukin-1 axis and cutaneous inflammation. *J Invest Dermatol* 1995;105:625-665.
97. Burger D, Dayer JM. Inhibitory cytokines and cytokine inhibitors. *Neurology* 1995;45:S39-S43.
98. Dinarello CA. The biological properties of interleukin-1. *Eur Cytokine Netw* 1994;5:517-531.
99. van der Meer JW, Vogels MT, Kullberg BJ. Interleukin-1 and related proinflammatory cytokines in the treatment of bacterial infections in neutropenic and non-neutropenic animals. *Biotherapy* 1994;7:161-167.
100. Mantovani A, Muzio M, Ghezzi P, Colotta F, Introna M. Negative regulators of the interleukin-1 system: receptor antagonists and a decoy receptor. *Int J Clin Lab Res* 1996;26:7-14.
101. Lennard AC. Interleukin-1 receptor antagonist. *Crit Rev Immunol* 1995;15:77-105.
102. Kopf M, Le Gros G, Coyle AJ, Kosco-Vilbois M, Brombacher F. Immune responses of IL-4, IL-5, IL-6 deficient mice. *Immunol Rev* 1995;148:45-69.
103. Puri RK. Structure and function of interleukin-4 and its receptor. *Cancer Treat Res* 1995;80:143-185.
104. Schroeder JT, Kagey-Sobotka A, MacGlashan DW, Lichtenstein LM. The interaction of cytokines with human basophils and mast cells. *Int Arch Allergy Immunol* 1995;107:79-81.
105. Tepper RI. The anti-tumor and proinflammatory actions of IL-4. *Res Immunol* 1993;144:633-637.
106. Ryan JJ, McReynolds LJ, Keegan A, et al. Growth and gene expression are predominantly controlled by distinct regions of the human IL-4 receptor. *Immunity* 1996;4:123-132.
107. Pernis A, Witthuhn B, Keegan AD, et al. Interleukin 4 signals through two related pathways. *Proc Natl Acad Sci USA* 1995;92:7971-7975.
108. Keegan AD, Johnston JA, Tortolani PJ, et al. Similarities and differences in signal transduction by interleukin 4 and interleukin 13: analysis of Janus kinase activation. *Proc Natl Acad Sci USA* 1995;92:7681-7685.
109. Sehgal PB. Interleukin-6-type cytokines in vivo: regulated bioavailability. *Proc Soc Exp Biol Med* 1996;213:238-247.
110. Biffi WL, Moore EE, Moore FA, Peterson VM. Interleukin-6 and the injured patient. Marker of injury or mediator of inflammation? *Ann Surg* 1996;224:647-664.
111. Baatout S. Interleukin-6 and megakaryocytopoiesis: an update. *Ann Hematol* 1996;73:157-162.
112. Paquet P, Pierard GE. Interleukin-6 and the skin. *Int Arch Allergy Immunol* 1996;109:308-317.
113. Luheshi G, Rothwell N. Cytokines and fever. *Int Arch Allergy Immunol* 1996;109:301-307.
114. Harada A, Mukaida N, Matsushima K. Interleukin 8 as a novel target for intervention therapy in acute inflammatory diseases. *Mol Med Today* 1996;2:482-489.
115. Hoch RC, Schraufstatter IU, Cochrane CG. In vivo, in vitro, and molecular aspects of interleukin-8 and the interleukin-8 receptors. *J Lab Clin Med* 1996;128:134-145.
116. Kunkel SL, Lukacs N, Streiter RM. Chemokines and their role in human disease. *Agents Actions Suppl* 1995;46:11-22.
117. Baggiolini M, Loetscher P, Moser B. Interleukin-8 and the chemokine family. *Int J Immunopharmacol* 1995;17:103-108.
118. Ben-Baruch A, Michiel DF, Oppenheim JJ. Signals and receptors involved in recruitment of inflammatory cells. *J Biol Chem* 1995;270:11703-11706.
119. Harada A, Sekido N, Akahoshi T, Wada T, Mukaida N, Matsushima K. Essential involvement of interleukin-8 (IL-8) in acute inflammation. *J Leukoc Biol* 1994;56:559-564.
120. Horuk R. The interleukin-8-receptor family: from chemokines to malaria. *Immunol Today* 1994;15:169-174.
121. Murphy PM, Tiffany HL. Cloning of a complementary DNA encoding a functional human interleukin-8 receptor. *Science* 1991;253:1280-1283.

122. Lynch DH. The role of FasL and TNF in the homeostatic regulation of immune responses. *Adv Exp Med Biol* 1996;406:135-138.
123. Barbara JA, van Ostade X, Lopez A. Tumour necrosis factor alpha (TNF-alpha): the good, the bad and the potentially very effective. *Immunol Cell Biol* 1996;74:434-443.
124. Riches DW, Chan ED, Winston BW. TNF-alpha-induced regulation and signalling in macrophages. *Immunobiology* 1996;195:477-490.
125. Douni E, Akassoglou K, Alexopoulos L, et al. Transgenic and knockout analyses of the role of TNF in immune regulation and disease pathogenesis. *J Inflamm* 1995;6:47:27-38.
126. Gruss HJ, Duyster J, Hermann F. Structural and biological features of the TNF receptor and TNF ligand superfamilies. *Ann Oncol* 1996;4:19-26.
127. Cosman D. A family of ligands for the TNF receptor superfamily. *Stem Cells* 1994;12:440-455.
128. Beutler B. TNF, immunity and inflammatory disease: lessons of the past decade. *J Invest Med* 1995;43:227-235.
129. Cleveland JL, Ihle JN. Contenders in the FasL/TNF death signalling. *Cell* 1995;81:479-482.
130. Ware CF, VanArsdale S, VanArsdale TL. Apoptosis mediated by the TNF-related cytokine and receptor families. *J Cell Biochem* 1996;60:47-55.
131. Baker SJ, Reddy EP. Transducers of life and death: TNF receptor superfamily and associated proteins. *Oncogene* 1996;12:1-9.
132. Young HA. Regulation of interferon-gamma gene expression. *J Interferon Cytokine Res* 1996;16:563-568.
133. Billiau A. Interferon-gamma: biology and role in pathogenesis. *Adv Immunol* 1996;62:61-130.
134. Young HA, Hardy KJ. Role of interferon-gamma in immune cell regulation. *J Leukoc Biol* 1995;58:373-381.
135. Gallin JI, Farber JM, Holland SM, Nutman TB. Interferon-gamma in the management of infectious diseases. *Ann Intern Med* 1995;123:216-224.
136. Appelberg R. Protective role of interferon gamma, tumor necrosis factor alpha and interleukin-6 in mycobacterium tuberculosis and M. avium infections. *Immunobiology* 1994;191:520-525.
137. Vilcek J, Oliveira JC. Recent progress in the elucidation of interferon-gamma actions: molecular biology and biological functions. *Int Arch Allergy Immunol* 1994;104:311-316.
138. Newport MJ, Huxley CM, Huston S, et al. A mutation in the interferon-gamma-receptor gene and susceptibility to mycobacterial infection. *N Engl J Med* 1996;335:1941-1949.
139. Holland SM, Gerstberger SM, Pitha-Rowe IF, et al. Interferon gamma receptor deficiency in disseminated mycobacterium avium complex (DMAC) infection. *J Invest Med* 1997;45:199A.
140. Brown CC, Malech HL, Gallin JI. Intravenous endotoxin recruits distinct subset of human neutrophils, defined by monoclonal antibody 3108, from bone marrow to the peripheral circulation. *Cell Immunol* 1989;123:294-306.
141. Zimmerli W, Gallin JI. Monocytes accumulate on Rebut skin window cover-slips but not in skin chamber fluid. *J Immunol Methods* 1987;96:11-17.
142. Kuhns DB, DeCarlo E, Hawk DM, Gallin JI. Dynamics of the cellular and humoral components of the inflammatory response elicited in skin blisters in human. *J Clin Invest* 1992;89:1734-1740.
143. Martich GD, Danner RL, Ceska M, Suffredini AF. Detection of interleukin-8 and tumor necrosis factor in normal humans after intravenous endotoxin: the effects of anti-inflammatory agents. *J Exp Med* 1991;173:1021-1024.
144. Kishimoto TK, Anderson DC. The role of integrins in inflammation. In: Gallin JI, Goldstein IM, Snyderman R, eds. *Inflammation*. 2nd ed. New York: Raven Press, 1992:353-406.
145. Kuhns DB, Long Priel DA, Gallin JI. Endotoxin and IL-1 hyporesponsiveness in a patient with recurrent bacterial infections. *J Immunol* 1997;158:3959-3964.
146. Haziot A, Tsuberi BZ, Goyert SM. Neutrophil CD14: biochemical properties and role in the secretion of tumor necrosis factor alpha in response to lipopolysaccharide. *J Immunol* 1993;150:5556-5565.
147. Frey EA, Miller DS, Jahr TG, et al. Soluble CD14 participates in the response of cells to lipopolysaccharide. *J Exp Med* 1992;176:1665-1671.
148. Muller JM, Loms Ziegler-Heitbrock HW, Bauerle PA. Nuclear factor kappa B, a mediator of lipopolysaccharide effects. *Immunobiology* 1993;187:233-256.
149. Smith JA. Neutrophils, host defense, and inflammation: a double-edged sword. *J Leukoc Biol* 1994;56:672-686.
150. Edwards SW. *Biochemistry and physiology of the neutrophil*. Cambridge: Cambridge University Press, 1994.
151. Domachowski JB, Malech HL. Phagocytes. In: Rich RR, Fleisher TA, Schwartz BD, Shearer WT, Strober W, eds. *Clinical immunology: principles and practice*. St. Louis: Mosby-Year Book, 1996:392-407.
152. Bainton DF. Developmental biology of neutrophils and eosinophils. In: Gallin JI, Goldstein IM, Snyderman R, eds. *Inflammation*. 2nd ed. New York: Raven Press, 1992:303-324.
153. Sievers EL, Dale DC. Non-malignant neutropenia. *Blood Rev* 1996;10:95-100.
154. Kim SK, Demetri GD. Chemotherapy and neutropenia. *Hematol Oncol Clin North Am* 1996;10:377-395.
155. Bernini JC. Diagnosis and management of chronic neutropenia during childhood. *Pediatr Clin North Am* 1996;43:773-792.
156. Welte K, Dale D. Pathophysiology and treatment of severe chronic neutropenia. *Ann Hematol* 1996;72:158-165.
157. Souid AK. Congenital cyclic neutropenia. *Clin Pediatr* 1995;34:151-155.
158. Cohen MS. Molecular events in the activation of human neutrophils for microbial killing. *Clin Infect Dis* 1994;18[Suppl 2]:S170-S179.
159. Downey GP, Fukushima T, Fialkow L, Waddell TK. Intracellular signaling in neutrophil priming and activation. *Semin Cell Biol* 1995;6:345-356.
160. Sengelov H. Complement receptors in neutrophils. *Crit Rev Immunol* 1995;15:107-131.
161. Edwards SW. Cell signalling by integrins and immunoglobulin receptors in primed neutrophils. *Trends Biochem Sci* 1995;20:362-367.
162. Murphy PM. The molecular biology of leukocyte chemoattractant receptors. *Annu Rev Immunol* 1994;12:593-633.
163. Wells TN, Lusti-Narasimhan M, Chung CW, et al. The molecular basis of selectivity between CC and CXC chemokines: the possibility of chemokine antagonists as anti-inflammatory agents. *Ann NY Acad Sci* 1996;796:245-256.
164. Hoch RC, Schraufstatter IU, Cochrane CG. In vivo, in vitro, and molecular aspects of interleukin-8 and the interleukin-8 receptors. *J Lab Clin Med* 1996;128:134-145.
165. Symon FA, Wardlaw AJ. Selectins and their counter receptors: a bitter sweet attraction. *Thorax* 1996;51:1155-1157.
166. Crockett-Torabi E, Fantone JC. The selectins: insights into selectin-induced intracellular signalling in leukocytes. *Immunol Res* 1995;14:237-251.
167. Kansas GS. Selectins and their ligands: current concepts and controversies. *Blood* 1996;88:3259-3287.
168. Vestweber D. Ligand-specificity of the selectins. *J Cell Biochem* 1996;61:585-591.
169. Tedder TF, Steeber DA, Chen A, Engel P. The selectins: vascular adhesion molecules. *FASEB J* 1995;9:866-873.
170. Rosen SD, Bertozzi CR. The selectins and their ligands. *Curr Opin Cell Biol* 1994;6:663-673.
171. McEver RP. Role of selectins in leukocyte adhesion to platelets and endothelium. *Ann NY Acad Sci* 1994;714:185-189.
172. McEver RP. Selectins. *Curr Opin Immunol* 1994;6:75-84.
173. Furie B, Furie BC. The molecular basis of platelet and endothelial cell interaction with neutrophils and monocytes: role of P-selectin and the P-selectin ligand PSGL-1. *Thromb Haemost* 1995;74:224-227.
174. Gille J, Swerlick RA. Integrins: role in cell adhesion and communication. *Ann NY Acad Sci* 1996;797:93-106.
175. Mazonne A, Ricevuti G. Leukocyte CD11/CD18 integrins: biological and clinical relevance. *Haematologica* 1995;80:161-175.
176. Edwards SW. Cell signalling by integrins and immunoglobulin receptors in primed neutrophils. *Trends Biochem Sci* 1995;20:362-367.
177. Luscinskas FW, Lawler J. Integrins as dynamic regulators of vascular function. *FASEB J* 1994;8:929-938.
178. Kishimoto TK, Rothlein R. Integrins, ICAMs and selectins: role and regulation of adhesion molecules in neutrophil recruitment to inflammatory sites. *Adv Pharmacol* 1994;25:117-169.
179. Patarroyo M. Adhesion molecules mediating recruitment of monocytes to inflamed tissue. *Immunobiology* 1994;191:474-477.
180. Carlos TM, Harlan JM. Leukocyte-endothelial adhesion molecules. *Blood* 1994;84:2068-2101.
181. van de Stolpe A, van der Saag PT. Intercellular adhesion molecule-1. *J Mol Med* 1996;74:13-33.
182. Anderson DC, Schmalstieg FC, Finegold MJ, et al. The severe and moderate phenotypes of heritable Mac-1, LFA-1 deficiency: their quantitative definition and relation to leukocyte dysfunction and clinical features. *J Infect Dis* 1988;152:668-689.
183. Anderson DC, Springer TA. Leukocyte adhesion deficiency and inherited defect in the Mac-1, LFA-1, and p150, 95 glycoproteins. *Ann Rev Med* 1987;175-194.
184. Kishimoto TK, Hollander N, Roberts TM, Anderson DC, Springer TA. Heterogeneous mutations in the beta subunit common to the LFA-1, Mac-1 and p150, 95 glycoproteins cause leukocyte adhesion deficiency. *Cell* 1987;50:193-202.
185. Etzioni A, Frydman M, Pollack S, et al. Brief report: recurrent severe infections caused by a novel leukocyte adhesion deficiency. *N Engl J Med* 1992;327:1789-1792.
186. Etzioni A, Phillips LM, Paulson JC, Harlan JM. Leukocyte adhesion deficiency (LAD) II. *Ciba Found Symp* 1995;189:51-58.
187. Brown SS. Structure and function of profilin. *Cell Motil Cytoskeleton* 1990;17:71-75.
188. Southwick FS, Stossel TP. Contractile proteins in leukocyte function. *Semin Hematol* 1984;30:305-310.
189. Stossel TP. The mechanical responses of white blood cells. In: Gallin JI, Goldstein IM, Snyderman R, eds. *Inflammation*. 2nd ed. New York: Raven Press, 1992:459-476.
190. Wright SD. Receptors for complement and the biology of phagocytosis. In: Gallin JI, Goldstein IM, Snyderman R, eds. *Inflammation*. 2nd ed. New York: Raven Press, 1992:477-496.
191. Allen LA, Aderem A. Mechanisms of phagocytosis. *Curr Opin Immunol* 1996;8:36-40.
192. Goldman MJ, Anderson GM, Stolzenberg ED, Kari UP, Zasloff M, Wilson JM. Human beta-defensin-1 is a salt-sensitive antibiotic in lung that is inactivated in cystic fibrosis. *Cell* 1997;88:553-560.
193. Lonnerdal B, Iyer S. Lactoferrin: molecular structure and biological function. *Annu Rev Nutr* 1995;15:93-110.
194. Nagle DL, Karim MA, Woolf EA, et al. Identification and mutation analysis of the complete gene for Chediak-Higashi syndrome. *Nat Genet* 1996;14:307-311.
195. Barbosa MD, Nguyen QA, Tchernev VT, et al. Identification of the homologous beige and Chediak-Higashi syndrome genes. *Nature* 1996;382:262-265.
196. Barrat FJ, Auloge L, Pastural E, et al. Genetic and physical mapping of the Ch-

- diak-Higashi syndrome on chromosome 1q42-43. *Am J Hum Genet* 1996;59:625-632.
197. Gallin JI. Neutrophil specific granule deficiency. *Annu Rev Med* 1985;36:263-274.
 198. Rosenberg HF, Gallin JI. Neutrophil specific granule deficiency includes eosinophils. *Blood* 1993;82:268-273.
 199. Lomax KJ, Gallin JI, Rotrosen D, et al. Selective defect in myeloid cell lactoferrin gene expression in neutrophil specific granule deficiency. *J Clin Invest* 1989;83:514-519.
 200. Parry MF, Root RK, Metcalf JA, Delaney KK, Kaplow LS, Richar WJ. Myeloperoxidase deficiency: prevalence and clinical significance. *Ann Intern Med* 1981;95:293-301.
 201. Nauseef WM, Brigham S, Cogley M. Hereditary myeloperoxidase deficiency due to a missense mutation of arginine 569 to tryptophan. *J Biol Chem* 1994;269:1212-1216.
 202. Nauseef WM, Cogley M, McCormack S. Effect of R569W missense mutation on the biosynthesis of myeloperoxidase. *J Biol Chem* 1996;271:9546-9549.
 203. DeLeo FR, Quinn MT. Assembly of the phagocyte NADPH oxidase: molecular interaction of the oxidase proteins. *J Leukoc Biol* 1996;60:677-691.
 204. Leusen JH, Verhoeven AJ, Roos D. Interactions between the components of the human NADPH oxidase: intrigues in the phox family. *J Lab Clin Med* 1996;128:461-476.
 205. Wientjes FB, Segal AW. NADPH oxidase and the respiratory burst. *Semin Cell Biol* 1995;6:357-365.
 206. Henderson LM, Chappel JB. NADPH oxidase of neutrophils. *Biochim Biophys Acta* 1996;1273:87-107.
 207. Segal AW. The NADPH oxidase and chronic granulomatous disease. *Mol Med Today* 1996;2:129-135.
 208. Jones OT. The regulation of superoxide production by the NADPH oxidase of neutrophils and other mammalian cells. *Bioessays* 1994;16:919-923.
 209. Bokoch GM. Regulation of the human neutrophil NADPH oxidase by the Rac GTP-binding proteins. *Curr Opin Cell Biol* 1994;6:212-218.
 210. Mardiny MM, Jackson SH, Spratt SK, Li F, Holland SM, Malech HL. Enhanced host defense after gene transfer in the murine p47-phox-deficient model of chronic granulomatous disease. *Blood* 1997;89:2268-2275.
 211. Lloyd AR, Oppenheim JJ. Poly's lament: the neglected role of the polymorphonuclear neutrophil in the afferent limb of the immune response. *Immunol Today* 1992;13:169-172.
 212. Granelli-Piperno A, Vassalli JD, Reich E. Secretion of plasminogen activator by human polymorphonuclear leukocytes. *J Exp Med* 1977;149:284-289.
 213. Shirafuji N, Matsuda S, Ogura H, et al. Granulocyte colony-stimulating factor stimulates human mature neutrophilic granulocytes to produce interferon-alpha. *Blood* 1990;75:17-19.
 214. Tiku K, Tiku ML, Skosey JL. Interleukin 1 production by human polymorphonuclear neutrophils. *J Immunol* 1986;136:3677-3685.
 215. Hughes V, Humphreys JM, Edwards SW. Protein synthesis is activated in primed neutrophils: a possible role in inflammation. *Biosci Rep* 1987;7:881-889.
 216. Kuhns DB, Gallin JI. Increased cell-associated IL-8 in human exudate and A23187-treated peripheral blood neutrophils. *J Immunol* 1995;154:6556-6662.
 217. van Furth R. Development and distribution of mononuclear phagocytes. In: Gallin JI, Goldstein IM, Snyderman R, eds. *Inflammation*. 2nd ed. New York: Raven Press, 1992:325-340.
 218. Adams DO, Hamilton TA. Macrophages as destructive cells in host defense. In: Gallin JI, Goldstein IM, Snyderman R, eds. *Inflammation*. 2nd ed. New York: Raven Press, 1992:637-662.
 219. Laskin DL, Pendino KJ. Macrophages and inflammatory mediators in tissue injury. *Annu Rev Pharmacol Toxicol* 1995;35:655-677.
 220. Cavaillon JM. Cytokines and macrophages. *Biomed Pharmacother* 1994;48:445-453.
 221. Thepen T, Kraal G, Holt PG. The role of alveolar macrophages in regulation of lung inflammation. *Ann NY Acad Sci* 1994;725:200-206.
 222. Seljelid R, Busund LT. The biology of macrophages: II. Inflammation and tumors. *Eur J Haematol* 1994;52:1-12.
 223. Nielsen BW, Mukaida N, Matsushima K, Kasahara T. Macrophages as producers of chemotactic proinflammatory cytokines. *Immunol Ser* 1994;60:131-142.
 224. Gleich GJ, Adolphson CR, Leiferman KM. Eosinophils. In: Gallin JI, Goldstein IM, Snyderman R, eds. *Inflammation*. 2nd ed. New York: Raven Press, 1992:663-700.
 225. Spry CJF. *Eosinophils: a comprehensive review and guide to the scientific and medical literature*. Oxford: Oxford University Press, 1988.
 226. Martin LB, Kita H, Leiferman KM, Gleich GJ. Eosinophils in allergy: role in disease, degranulation and cytokines. *Int Arch Allergy Immunol* 1996;109:207-215.
 227. Makino S, Fukuda T. *Eosinophils: biological and clinical aspects*. Boca Raton, FL: CRC Press, 1993.
 228. Wardlaw AJ. Eosinophils in the 1990s: new perspectives on their role in health and disease. *Postgrad Med J* 1994;70:536-552.
 229. Makino S, Fukuda T. Eosinophils and allergy in asthma. *Allergy Proc* 1995;16:13-21.
 230. Seminario MC, Gleich GJ. The role of eosinophils in the pathogenesis of asthma. *Curr Opin Immunol* 1994;6:860-864.
 231. Reed CE. The importance of eosinophils in the immunology of asthma and allergic disease. *Ann Allergy* 1994;72:376-380.
 232. Weksler BB. Platelets. In: Gallin JI, Goldstein IM, Snyderman R, eds. *Inflammation*. 2nd ed. New York: Raven Press, 1992:727-746.
 233. de Sousa JR, Palma-Carlos AG. Platelets and hypersensitivity. *J Invest Allergol Clin Immunol* 1995;5:12-17.
 234. Herd CM, Page CP. Pulmonary immune cells in health and disease: platelets. *Eur Respir J* 1994;7:1145-1160.
 235. Nash GB. Adhesion between neutrophils and platelets: a modulator of thrombotic and inflammatory events? *Thromb Res* 1994;74:S3-S11.
 236. Tonnel AB, Gosset P, Molet S, Tillie-Leblond I, Jeannin P, Joseph M. Interactions between endothelial cells and effector cells in allergic inflammation. *Ann NY Acad Sci* 1996;796:9-20.
 237. Kubes P, Granger DN. Leukocyte-endothelial cell interactions evoked by mast cells. *Cardiovasc Res* 1996;32:699-708.
 238. Malik AB, Lo SK. Vascular endothelial adhesion molecules and tissue inflammation. *Pharmacol Rev* 1996;48:213-229.
 239. Lusinskas FW, Gimbrone MA Jr. Endothelial-dependent mechanisms in chronic inflammatory leukocyte recruitment. *Annu Rev Med* 1996;47:413-421.
 240. Garcia JG, Pavalko FM, Patterson CE. Vascular endothelial cell activation and permeability responses to thrombin. *Blood Coagul Fibrinolysis* 1995;6:609-626.
 241. Chen CC, Manning AM. Transcriptional regulation of endothelial cell adhesion molecules: a dominant role for NF-kappa B. *Agents Actions Suppl* 1995;47:135-141.
 242. Bevilacqua MP, Nelson RM, Mannori G, Cecconi O. Endothelial-leukocyte adhesion molecules in human disease. *Annu Rev Med* 1994;45:361-378.
 243. Vane JR, Botting RM. Mediators from the endothelial cell and their participation in inflammation. *Int J Tissue React* 1994;16:19-49.
 244. Varani J, Ward PA. Mechanisms of neutrophil-dependent and neutrophil-independent endothelial cell injury. *Biol Signals* 1994;3:1-14.
 245. Korthuis RJ, Anderson DC, Granger DN. Role of neutrophil-endothelial cell adhesion in inflammatory disorders. *J Crit Care* 1994;9:47-71.
 246. Granger DN, Kubes P. The microcirculation and inflammation. Modulation of leukocyte-endothelial cell adhesion. *J Leukoc Biol* 1994;55:662-675.
 247. Benedict CR, Pakala R, Willerson JT. Endothelial-dependent procoagulant and anti-coagulant mechanisms. Recent advances in understanding. *Tex Heart Inst J* 1994;21:86-90.
 248. Jarrett EEE, Bazin H. Elevation of total serum IgE in rats following helminth parasite infection. *Nature* 1974;251:541-543.
 249. Jarrett E. Stimuli for the production and control of IgE in rats. *Immunol Rev* 1978;41:52-76.
 250. Hussain R, Ottesen EA. IgE responses in human filariasis. 3. Specificities of IgE and IgG antibodies compared by immunoblot analysis. *J Immunol* 1985;135:1415-1420.
 251. Anderson GP. Resolution of chronic inflammation by therapeutic induction of apoptosis. *Trends Pharmacol Sci* 1996;17:438-442.
 252. Savill J, Haslett C. Granulocyte clearance by apoptosis in the resolution of inflammation. *Semin Cell Biol* 1995;6:385-393.
 253. Bellamy CO, Malcolmson RD, Harrision DJ, Wyllie AH. Cell death in health and disease: the biology and regulation of apoptosis. *Semin Cancer Biol* 1995;6:3-16.
 254. Savill J. Apoptosis in disease. *Eur J Clin Invest* 1994;24:715-723.
 255. Haslett C, Savill JS, Whyte MK, Stern M, Dransfield I, Meagher LC. Granulocyte apoptosis and the control of inflammation. *Philos Trans R Soc London B Biol Sci* 1994;345:327-333.
 256. Simon HU, Yousefi S, Blaser K. Tyrosine phosphorylation regulated activation and inhibition of apoptosis in human eosinophils and neutrophils. *Int Arch Allergy Immunol* 1995;107:338-339.
 257. Liles WC, Klebanoff SJ. Regulation of apoptosis in neutrophils—Fas track to death? *J Immunol* 1995;155:3289-3291.
 258. Cox G. IL-10 enhances resolution of pulmonary inflammation in vivo by promoting apoptosis of neutrophils. *Am J Physiol* 1996;27:L566-L571.
 259. Gelrud AK, Carper HT, Mandell GL. Interaction of tumor necrosis factor-alpha and granulocyte colony-stimulating factor on neutrophil apoptosis, receptor expression, and bactericidal function. *Proc Assoc Am Physicians* 1996;108:455-456.
 260. Gon S, Gatanaga T, Sendo F. Involvement of two types of TNF receptor in TNF-alpha induced neutrophil apoptosis. *Microbiol Immunol* 1996;40:463-465.
 261. Hebert MJ, Takano T, Holthofer H, Brady HR. Sequential morphologic events during apoptosis of human neutrophils. Modulation by lipoxygenase-derived eicosanoids. *J Immunol* 1996;157:3105-3115.
 262. Oishi K, Machida K. Inhibition of neutrophil apoptosis by antioxidants in culture medium. *Scand J Immunol* 1997;45:21-27.
 263. Abramson SL, Gallin JI. Interleukin-4 inhibits superoxide production by human mononuclear phagocytes. *J Immunol* 1990;144:625-630.
 264. Lawrence DA. Transforming growth factor-beta: a general review. *Eur Cytokine Netw* 1996;7:363-374.
 265. Kolodziejczyk SM, Hall BK. Signal transduction and TGF-beta superfamily receptors. *Biochem Cell Biol* 1996;74:299-314.
 266. Markowitz SD, Roberts AB. Tumor suppressor activity of the TGF-beta pathway in human cancers. *Cytokine Growth Factor Rev* 1996;7:93-102.
 267. Kingsley DM. The TGF-beta superfamily: new members, new receptors, and new genetic tests of function in different organisms. *Genes Dev* 1994;8:133-146.
 268. Serra R, Moses HL. Tumor suppressor genes in the TGF-beta signaling pathway? *Nat Med* 1996;2:390-391.
 269. Derynck R. TGF-beta-receptor-mediated signaling. *Trends Biochem Sci* 1994;19:548-553.
 270. Letterio JJ, Geiser AG, Kulkarni AB, et al. Autoimmunity associated with TGF-beta1-deficiency in mice is dependent on MHC class II antigen expression. *J Clin Invest* 1996;98:2109-2119.

271. Dang H, Geiser AG, Letterio JJ, et al. SLE-like autoantibodies and Sjogren's syndrome-like lymphoproliferation in TGF-beta knockout mice. *J Immunol* 1995; 155:3205-3212.
272. Moore KW, O Garra A, deWaal Malefyt R, Viera P, Mosmann TR. Interleukin-10. *Annu Rev Immunol* 1993;11:165-190.
273. Hoiden I, Moller G. CD8+ cells are the main producers of IL-10 and IFN gamma after superantigen stimulation. *Scand J Immunol* 1996;44:501-505.
274. Klein SC, Kube D, Abts H, Diehl V, Tesch H. Promotion of IL-8, IL-10, TNF-alpha and TNF-beta production by EBV infection. *Leuk Res* 1996;20:633-636.
275. Schuitemaker H. IL-4 and IL-10 as potent inhibitors of HIV-1 replication in macrophages in vitro. A role for cytokines in the in vivo virus host range? *Res Immunol* 1994;145:588-592.
276. Mosmann TR. Properties and functions of interleukin 10. *Adv Immunol* 1994;56: 1-26.
277. McKenzie AN, Culpepper JA, de Waal Malefyt R, et al. Interleukin 13, a T-cell-derived cytokine that regulated human monocyte and B-cell function. *Proc Natl Acad Sci USA* 1993;90:3735-3739.
278. de Vries JE, Punnonen J, Cocks BG, de Waal Malefyt R, Aversa G. Regulation of the human IgE response by IL-4 and IL-13. *Res Immunol* 1993;144:597-601.
279. Mossman T. Cytokines and immune regulation. In: Rich RR, Fleisher TA, Schwartz BD, Shearer WT, Strober W, eds. *Clinical immunology: principles and practice*. St. Louis: Mosby-Year Book, 1996:217-230.
280. Sternberg EM. Neuroendocrine factors in susceptibility to inflammatory disease: focus on the hypothalamic-pituitary-adrenal axis. *Horm Res* 1995;43: 159-161.
281. Buckingham JC, Loxley HD, Christian HC, Philip JG. Activation of the HPA axis by immune insults: roles and interactions of cytokines, eicosanoids and glucocorticoids. *Pharmacol Biochem Behav* 1996;54:285-298.
282. Derijk R, Sternberg EM. Corticosteroid action and neuroendocrine-immune interactions. *Ann NY Acad Sci* 1994;746:33-41.
283. Cunningham ET, de Souza EB. Interleukin-1 receptors in the brain and endocrine system. *Immunol Today* 1992;14:171-173.
284. Munck A, Guyre PM. Glucocorticoids and immune function. In: Ader R, Felton DL, Cohen N, eds. *Psychoneuroimmunology*. New York: Academic Press, 1991: 447-474.
285. Martin P. Wound healing—aiming for perfect skin regeneration. *Science* 1997; 276:75-81.
286. Davidson JM. Wound repair. In: Gallin JI, Goldstein IM, Snyderman R, eds. *Inflammation*. 2nd ed. New York: Raven Press, 1992:809-820.
287. Folkman J, Brem H. Angiogenesis and inflammation. In: Gallin JI, Goldstein IM, Snyderman R, eds. *Inflammation*. 2nd ed. New York: Raven Press, 1992: 821-840.
288. Strieter RM, Polverini PJ, Arenberg DA, Kunkel SL. The role of CXC chemokines as regulators of angiogenesis. *Shock* 1995;4:155-160.
289. Diaz-Flores L, Gutierrez R, Varela H. Angiogenesis: an update. *Histol Histopathol* 1994;9:807-843.
290. Boros DL. The role of cytokines in the formation of the schistosome egg granuloma. *Immunobiology* 1994;191:441-450.
291. Wynn TA, Cheever AW. Cytokine regulation of granuloma formation in schistosomiasis. *Curr Opin Immunol* 1995;7:505-511.
292. Staderker M. The shrinking schistosomal egg granuloma: how accessory cells control T cell-mediated pathology. *Exp Parasitol* 1994;79:198-201.
293. Mornex JF, Leroux C, Greenland T, Ecochard D. From granuloma to fibrosis in interstitial lung diseases: molecular and cellular interactions. *Eur Respir J* 1994; 7:779-785.
294. Kunkel SL, Lukacs NW, Strieter RM, Chensue SW. Th1 and Th2 responses regulate experimental lung granuloma development. *Sarcoidosis Vasc Diffuse Lung Dis* 1996;13:120-128.
295. Hansch HC, Smith DA, Mielke ME, Hahn H, Bancroft GJ, Ehlers S. Mechanisms of granuloma formation in murine *Mycobacterium avium* infection: the contribution of CD4+ T cells. *Int Immunol* 1996;8:1299-1310.
296. Chensue SW, Warmington KS, Ruth JH, Lincoln P, Kunkel SL. Cytokine function during mycobacterial and schistosomal antigen-induced pulmonary granuloma formation. Local and regional participation of IFN-gamma, IL-10, and TNF. *J Immunol* 1995;154:5969-5976.
297. Barnes PJ, Karin M. Nuclear factor kappa-B—a pivotal transcription factor in chronic inflammatory diseases. *N Engl J Med* 1997;336:1066-1071.
298. Barnes PJ, Adcock I. Anti-inflammatory actions of steroids: molecular mechanisms. *Trends Pharmacol Sci* 1993;14:436-441.
299. Saatcioglu F, Claret FX, Karin M. Negative transcriptional regulation by nuclear receptors. *Semin Cancer Biol* 1994;5:347-359.
300. Rosenberg HF, Gallin JI. Neutrophils. In: Frank MM, Austen KF, Claman HN, Unanue ER, eds. *Samter's immunologic diseases*. 5th ed. Boston: Little, Brown and Company, 1995:247-250.
301. Leto TL, Adams AG, deMendez I. Assembly of the phagocyte NADPH oxidase: binding of Src homology 3 domains to proline-rich targets. *Proc Natl Acad Sci USA* 1994;91:10650-10654.
302. deMendez I, Garrett MC, Adams AG, Leto TL. The role of p67-phox SH3 domains in assembly of the NADPH oxidase system. *J Biol Chem* 1994;269: 16326-16332.
303. Sathymoorthy M, deMendez I, Adams AG, Leto TL. P40 phox down-regulated NADPH oxidase activity through interactions with its SH3 domain. *J Biol Chem* 1997;272:9141-9146.
304. Jackson SH, Gallin JI, Holland SM. The p47 phox mouse knockout model of chronic granulomatous disease. *J Exp Med* 1995;182:751-758.
305. Marks AR. Cellular functions of immunophilins. *Physiol Rev* 1996;76:631-649.
306. Yocum DE. Cyclosporine, FK-506, rapamycin and other immunomodulators. *Rheum Dis Clin North Am* 1996;22:133-154.
307. Lee JJ, Canafax DM. Cyclosporine pharmacology. *Transplant Proc* 1996;28: 2156-2158.
308. Khanna A, Li B, Sharma VK, Suthanthiran M. Immunoregulatory and fibrogenic activities of cyclosporine: a unifying hypothesis based on transforming growth factor-beta expression. *Transplant Proc* 1996;28:2015-2018.
309. Stiller CR. An overview of the first decade of cyclosporine. *Transplant Proc* 1996;28:2005-2012.
310. High KP. The antimicrobial activities of cyclosporine, FK-506 and rapamycin. *Transplantation* 1994;57:1689-1700.
311. Emery P, van de Putte LB. New perspectives on controlling progression of severe rheumatoid arthritis—a summary of the potential role of cyclosporin. *Br J Rheumatol* 1996;35[Suppl 2]:1-3.
312. Yocum DE. Combination therapy with cyclosporin in rheumatoid arthritis. *Br J Rheumatol* 1996;35[Suppl 2]:19-23.
313. Forre O. Cyclosporine in rheumatoid arthritis: an overview. *Clin Rheumatol* 1995;14[Suppl 2]:33-36.
314. Yocum DE, Torley H. Cyclosporine in rheumatoid arthritis. *Rheum Dis Clin North Am* 1995;21:835-844.
316. Weckmann AL, Alcocer-Varela J. Cytokine inhibitors in autoimmune disease. *Semin Arthritis Rheum* 1996;26:539-557.
317. Feldmann M, Brennan FM, Williams RO, Elliott MJ, Maini RN. Cytokine expression and networks in rheumatoid arthritis: rationale for anti-TNF alpha therapy and its mechanism of action. *J Inflamm* 1995;6:47:90-96.
318. Sasayama S, Matsumori A, Vesnarione. A potential cytokine inhibitor. *J Card Fail* 1996;2:251-258.
319. Maini RN. A perspective on anti-cytokine and anti-T cell-directed therapies in rheumatoid arthritis. *Clin Exp Rheumatol* 1995;13[Suppl 12]:S35-S40.
320. Bartalena L, Marcocci C, Pinchera A. cytokine antagonists: new ideas for the management of Graves' ophthalmopathy. *J Clin Endocrinol Metab* 1996;81: 446-448.
321. Ferrara JL. Cytokine inhibitors and graft-versus-host disease. *Ann NY Acad Sci* 1995;770:227-236.
322. Richardson CE, Emery P. Innovative treatment approaches for rheumatoid arthritis. New cyclo-oxygenase and cytokine inhibitors. *Baillieres Clin Rheumatol* 1995;9:731-758.
323. Elliott MJ, Maini RN. Anti-cytokine therapy in rheumatoid arthritis. *Baillieres Clin Rheumatol* 1995;9:633-652.
324. Schwiebert LA, Beck LA, Stellato C, Bickel CA, Bochner BS, Schleimer RP. Glucocorticosteroid inhibition of cytokine production: relevance to antiallergic actions. *J Allergy Clin Immunol* 1996;97:143-152.
325. Feldmann M, Brennan FM, Elliott MJ, Williams RO, Maini RN. TNF-alpha is an effective therapeutic target for rheumatoid arthritis. *Ann NY Acad Sci* 1995; 766:272-278.
326. Border WA, Noble NA, Ketteler M. TGF-beta: a cytokine mediator of glomerulosclerosis and a target for therapeutic intervention. *Kidney Int Suppl* 1995;49: S59-S61.
327. Devos R, Plaetinck G, Cornelis S, Guisez Y, van der Heyden J, Tavernier J. Interleukin-5 and its receptor: a drug target for eosinophilia associated with chronic allergic disease. *J Leukoc Biol* 1995;57:813-819.
328. Firestein GS. Cytokine networks in rheumatoid arthritis: implication for therapy. *Agents Actions Suppl* 1995;47:37-51.
329. Klein B, Brailly H. Cytokine-binding proteins: stimulating antagonists. *Immunol Today* 1995;16:216-220.
330. Tartour E, Lee RS, Fridman WH. Anti-cytokines: promising tools for diagnosis and immunotherapy. *Biomed Pharmacother* 1994;48:417-424.
331. Enayati P, Fong Y. Cytokine neutralizing strategies in experimental sepsis. *Prog Clin Biol Res* 1994;388:295-306.
332. Kinoshita T. Protection of host from its own complement by membrane-bound complement inhibitors: C3 convertase inhibitors vs. membrane attack complex inhibitors. *Res Immunol* 1996;147:100-103.
333. Mouthon L, Kaveri SV, Spalter SH, et al. Mechanisms of action of intravenous immune globulin in immune-mediated diseases. *Clin Exp Immunol* 1996; 104[Suppl 1]:3-9.
334. Ryan US. Complement inhibitory therapeutics and xenotransplantation. *Nat Med* 1995;1:967-968.
335. Smiley JD, Talbert MG. Southwestern Internal Medicine Conference: high-dose intravenous gamma globulin therapy: how does it work? *Am J Med Sci* 1995;309:295-303.
336. Corral LG, Muller GW, Moreira AL, et al. Selection of novel analogs of thalidomide with enhanced tumor necrosis factor alpha inhibitory activity. *Mol Med* 1996;2:506-515.
337. Wang P, Wu P, Siegel MI, Egan RW, Billah MM. Interleukin (IL)-10 inhibits nuclear factor kappa B (NF kappa B) activation in human monocytes: IL-10 and IL-4 suppress cytokine synthesis by different mechanisms. *J Biol Chem* 1995; 270:9558-9563.
338. Pahl HL, Krauss B, Schulze-Osthoff K, et al. The immunosuppressive fungal metabolite gliotoxin specifically inhibits transcription factor NF-kappa B. *J Exp Med* 1996;183:1829-1834.